

gr-qc/0408080

LPT-ORSAY-04-64

UFIFT-QG-04-5

t04/103

Two Loop Scalar Self-Mass during Inflation

T. Brunier*

Service de Physique Théorique, CEA/Saclay, 91191 Gif-sur-Yvette FRANCE

V. K. Onemli†

Laboratoire de Physique Théorique, Université Paris XI, Bât. 210, 91405 Orsay FRANCE

R. P. Woodard‡

*Department of Physics, University of Florida, Gainesville, FL 32611 USA***ABSTRACT**

We work in the locally de Sitter background of an inflating universe and consider a massless, minimally coupled scalar with a quartic self-interaction. We use dimensional regularization to compute the fully renormalized scalar self-mass-squared at one and two loop order for a state which is released in Bunch-Davies vacuum at $t = 0$. Although the field strength and coupling constant renormalizations are identical to those of flat space, the geometry induces a non-zero mass renormalization. The finite part also shows a sort of growing mass that competes with the classical force in eventually turning off this system's super-acceleration.

PACS numbers: 98.80.Cq, 4.62.+v

* e-mail: brunier@spht.saclay.cea.fr

† e-mail: Vakif.Onemli@th.u-psud.fr

‡ e-mail: woodard@phys.ufl.edu

I. INTRODUCTION

For studying quantum fields which are spectators during inflation it is convenient to model the background geometry using a portion of the full de Sitter manifold known as the open coordinate patch. If the D -dimensional cosmological constant is $\Lambda \equiv (D-1)H^2$, the invariant element can be expressed, in either conformal or co-moving coordinates, as follows:

$$ds^2 = a^2(-d\eta^2 + d\vec{x} \cdot d\vec{x}) = -dt^2 + a^2 d\vec{x} \cdot d\vec{x} \quad \text{where} \quad a(\eta) = -\frac{1}{H\eta} = e^{Ht} . \quad (1)$$

One might expect that the inflationary expansion of this spacetime makes quantum effects stronger by allowing virtual particles to persist longer than in flat space. Indeed, it is simple to see that any sufficiently long wavelength (wavelength $> 1/H$) virtual particle which is massless on the Hubble scale can exist forever [1]. However, one must also consider the *rate* at which virtual particles emerge from the vacuum. Classical conformal invariance causes this rate (per unit co-moving time) to fall like $1/a$, so any long wave length virtual particles which emerge become real, but very few emerge [1]. To get enhanced quantum effects during inflation therefore requires quanta which are effectively massless and also not classically conformally invariant.

A fascinating example of enhanced quantum effects occurs when a massless minimally coupled scalar is endowed with a quartic self-interaction,

$$\mathcal{L} = -\frac{1}{2}\partial_\mu\varphi\partial_\nu\varphi g^{\mu\nu}\sqrt{-g} - \frac{\lambda}{4!}\varphi^4\sqrt{-g} + \text{counterterms} . \quad (2)$$

If this system is released at $t = 0$ ($\eta = -1/H$) in free Bunch-Davies vacuum the fully renormalized energy density and pressure are [2, 3],

$$\rho_{\text{ren}} = \frac{\Lambda}{8\pi G} + \frac{\lambda H^4}{2^6\pi^4} \left\{ \frac{1}{2} \ln^2(a) + \frac{2}{9}a^{-3} - \frac{1}{2} \sum_{n=1}^{\infty} \frac{n+2}{(n+1)^2} a^{-n-1} \right\} + O(\lambda^2) , \quad (3)$$

$$p_{\text{ren}} = -\frac{\Lambda}{8\pi G} - \frac{\lambda H^4}{2^6\pi^4} \left\{ \frac{1}{2} \ln^2(a) + \frac{1}{3} \ln(a) + \frac{1}{6} \sum_{n=1}^{\infty} \frac{n^2-4}{(n+1)^2} a^{-n-1} \right\} + O(\lambda^2) . \quad (4)$$

Hence their sum violates the weak energy condition on cosmological scales,

$$\rho_{\text{ren}} + p_{\text{ren}} = \frac{\lambda H^4}{2^6\pi^4} \left\{ -\frac{1}{3} \ln(a) + \frac{2}{9}a^{-3} - \frac{1}{6} \sum_{n=1}^{\infty} \frac{n+2}{n+1} a^{-n-1} \right\} + O(\lambda^2) . \quad (5)$$

The physics behind this surprising result is that inflationary particle production causes the scalar to undergo a random walk such that its average distance from the minimum of

the φ^4 potential increases without bound in the free theory [4–6],

$$\langle \Omega | \varphi^2(x) | \Omega \rangle_0 = (\text{Divergent Constant}) + \frac{H^2}{4\pi^2} \ln(a) . \quad (6)$$

After the ultraviolet divergence is removed by renormalization the effect is to drive the scalar up its potential, which increases the vacuum energy and leads to (3) and (4). The process is self-limiting for two reasons. First, as the scalar rises up its potential the classical φ^3 force pushes it back down. Second, the φ^2 curvature associated with being away from $\varphi = 0$ acts like a positive mass-squared to reduce the inflationary particle production responsible for pushing the scalar away from $\varphi = 0$.

This model is of more than passing interest because the supernova acceleration data are consistent with an evolving dark energy equation of state whose current value is less than minus one [7–18]. Such a thing is difficult to explain with classical models on account of the problem with stability [19–83]. Therefore one is lead naturally to a self-limiting quantum effect in a classically stable theory such as (2).

Although this model gives a value for $w + 1$ which is far too small to explain the observations [3], it does serve to establish that quantum effects can induce a temporary phase of super-acceleration. Further, it does so in a wonderfully simple setting for which every issue can be settled by explicit computation. To check stability one first computes the self-mass-squared $M^2(x; x')$ and then studies how it modifies the effective field equations at linearized order,

$$\square \varphi(x) - \int d^4x' M^2(x; x') \varphi(x') = 0 . \quad (7)$$

That first step is the object of this paper. Just as in flat space one must go to two loop order to see corrections to the derivative terms, so we will compute the fully renormalized self-mass-squared at one and two loop orders. In a later work we will employ the result to study the behavior of linearized perturbations in the quantum-corrected field equations.

In Section II we explain the Schwinger-Keldysh formalism that one must use for cosmological settings in which the state is specified at a finite time rather than in the asymptotic past and future. We also derive the Feynman rules. The one loop counterterms are computed in Section III. Sections IV–VI treat the various two loop contributions to the self-mass-squared. In Section VII we give the fully renormalized results. Implications are discussed in Section VIII. Section IX is an appendix devoted to various technical results used throughout.

II. FEYNMAN RULES IN THE SCHWINGER-KELDYSH FORMALISM

The Schwinger-Keldysh formalism is a technique that makes computing expectation values almost as simple as the Feynman rules do for computing in-out matrix elements [84, 85]. To sketch the derivation, consider a real scalar field, $\phi(x)$ whose Lagrangian (not Lagrangian density) at time t is $L[\phi(t)]$. Most quantum field theorists are familiar with the functional integral expression for the matrix element of an operator $\mathcal{O}_1[\phi]$ between states whose wave functionals are given at a starting time s and a last time ℓ ,

$$\langle \Phi | T^* (\mathcal{O}_1[\phi]) | \Psi \rangle = \int [d\phi] \mathcal{O}_1[\phi] \Phi^*[\phi(\ell)] e^{i \int_s^\ell dt L[\phi(t)]} \Psi[\phi(s)] . \quad (8)$$

The T^* -ordering symbol in the matrix element indicates that the operator $\mathcal{O}_1[\phi]$ is time-ordered, except that any derivatives are taken *outside* the time-ordering. We can use (8) to obtain a similar expression for the matrix element of the *anti*-time-ordered product of some operator $\mathcal{O}_2[\phi]$ in the presence of the reversed states,

$$\langle \Psi | \overline{T}^* (\mathcal{O}_2[\phi]) | \Phi \rangle = \langle \Phi | T^* (\mathcal{O}_2^\dagger[\phi]) | \Psi \rangle^* \quad (9)$$

$$= \int [d\phi] \mathcal{O}_2[\phi] \Phi[\phi(\ell)] e^{-i \int_s^\ell dt L[\phi(t)]} \Psi^*[\phi(s)] . \quad (10)$$

Now note that summing over a complete set of states Φ gives a delta functional,

$$\sum_{\Phi} \Phi[\phi_-(\ell)] \Phi^*[\phi_+(\ell)] = \delta[\phi_-(\ell) - \phi_+(\ell)] . \quad (11)$$

Taking the product of (8) and (10), and using (11), we obtain a functional integral expression for the expectation value of any anti-time-ordered operator $\mathcal{O}_2$ multiplied by any time-ordered operator $\mathcal{O}_1$,

$$\begin{aligned} \langle \Psi | \overline{T}^* (\mathcal{O}_2[\phi]) T^* (\mathcal{O}_1[\phi]) | \Psi \rangle &= \int [d\phi_+][d\phi_-] \delta[\phi_-(\ell) - \phi_+(\ell)] \\ &\times \mathcal{O}_2[\phi_-] \mathcal{O}_1[\phi_+] \Psi^*[\phi_-(s)] e^{i \int_s^\ell dt \left\{ L[\phi_+(t)] - L[\phi_-(t)] \right\}} \Psi[\phi_+(s)] . \end{aligned} \quad (12)$$

This is the fundamental relation between the canonical operator formalism and the functional integral formalism in the Schwinger-Keldysh formalism.

The Feynman rules follow from (12) in close analogy to those for in-out matrix elements. Because the same field is represented by two different dummy functional variables, $\phi_\pm(x)$, the endpoints of lines carry a $\pm$ polarity. External lines associated with the operator $\mathcal{O}_2[\phi]$

have $-$ polarity whereas those associated with the operator $\mathcal{O}_1[\phi]$ have $+$ polarity. Interaction vertices are either all $+$ or all $-$. Vertices with $+$ polarity are the same as in the usual Feynman rules whereas vertices with the $-$ polarity have an additional minus sign. Propagators can be $++$, $-+$, $+-$ and $--$.

The four propagators can be read off from the fundamental relation (12) when the free Lagrangian is substituted for the full one. It is useful to denote canonical expectation values in the free theory with a subscript 0. With this convention we see that the $++$ propagator is just the ordinary Feynman propagator,

$$i\Delta_{++}(x; x') = \langle \Omega | T(\phi(x)\phi(x')) | \Omega \rangle_0 = i\Delta(x; x'). \quad (13)$$

The other cases are simple to read off and to relate to the Feynman propagator,

$$i\Delta_{-+}(x; x') = \langle \Omega | \phi(x)\phi(x') | \Omega \rangle_0 = \theta(t-t')i\Delta(x; x') + \theta(t'-t)[i\Delta(x; x')]^*, \quad (14)$$

$$i\Delta_{+-}(x; x') = \langle \Omega | \phi(x')\phi(x) | \Omega \rangle_0 = \theta(t-t')[i\Delta(x; x')]^* + \theta(t'-t)i\Delta(x; x'), \quad (15)$$

$$i\Delta_{--}(x; x') = \langle \Omega | \overline{T}(\phi(x)\phi(x')) | \Omega \rangle_0 = [i\Delta(x; x')]^*. \quad (16)$$

Therefore we can get the four propagators of the Schwinger-Keldysh formalism from the Feynman propagator once that is known.

Quantum field theory has much the same structure in the Schwinger-Keldysh formalism as for the familiar in-out formalism. In particular, one-particle-irreducible (1PI) diagrams mean the same thing and play the same fundamental role. Because each external line can take either polarity there are 2^N variations for each N -point function of the in-out formalism. They generate an effective action that depends upon background fields $\phi_+(x)$ and $\phi_-(x)$. For example, there are four Schwinger-Keldysh 2-point 1PI functions, $M_{\pm\pm}^2(x; x')$. The $++$ one is the same as the in-out self-mass-squared and the others are related as the propagators,

$$-iM_{--}^2(x; x') = \left(-iM_{++}^2(x; x')\right)^* \quad , \quad -iM_{-+}^2(x; x') = \left(-iM_{+-}^2(x; x')\right)^*. \quad (17)$$

The various self-mass-squared's enter the effective action as follows [85],

$$\Gamma[\phi_+, \phi_-] = S[\phi_+] - S[\phi_-] - \frac{1}{2} \int d^D x \int d^D x' \left\{ \begin{aligned} &\phi_+(x)M_{++}^2(x; x')\phi_+(x') + \phi_+(x)M_{+-}^2(x; x')\phi_-(x') \\ &+ \phi_-(x)M_{-+}^2(x; x')\phi_+(x') + \phi_-(x)M_{--}^2(x; x')\phi_-(x') \end{aligned} \right\} + O(\phi_{\pm}^3). \quad (18)$$

Note that since counterterms derive from the classical actions — $S[\phi_+]$ and $-S[\phi_-]$ — there can be no primitive divergences in 1PI diagrams of mixed polarity.

The effective field equations of the Schwinger-Keldysh formalism are obtained by varying with respect to either background field and then setting the two fields equal [85],

$$\left. \frac{\delta \Gamma[\phi_+, \phi_-]}{\delta \phi_+(x)} \right|_{\phi_\pm = \phi} = \frac{\delta S[\phi]}{\delta \phi(x)} - \int d^D x' \{ M_{++}^2(x; x') + M_{+-}^2(x; x') \} \phi(x') + O(\phi^2) = 0. \quad (19)$$

Just as in-out matrix elements of the field obey the in-out effective field equations, so expectation values of the field obey the Schwinger-Keldysh effective field equations. Unlike the in-out effective field equations, the Schwinger-Keldysh effective field equations are causal and purely real. That is, although the equations are generally not local, there is no net dependence at point x^μ from fields at any point x'^μ which lies outside the past light-cone of x^μ . Hence the initial value problem is well defined. Also, if the initial value data are real, the solution remains real.

Up to this point the discussion has been general. We begin the process of deriving the Feynman rules of our model by expressing its Lagrangian (2) in terms of the unrenormalized field ϕ and bare parameters m_0^2 and λ_0 ,

$$\mathcal{L} = -\frac{1}{2} \partial_\mu \phi \partial_\nu \phi g^{\mu\nu} \sqrt{-g} - \frac{1}{2} m_0^2 \phi^2 \sqrt{-g} - \frac{1}{4!} \lambda_0 \phi^4 \sqrt{-g}. \quad (20)$$

Specializing to the (D-dimensional) locally de Sitter geometry in conformal coordinates (1), and introducing the renormalized field $\varphi \equiv \phi/\sqrt{Z}$ gives,

$$\mathcal{L} = -\frac{1}{2} Z \partial_\mu \varphi \partial_\nu \varphi \eta^{\mu\nu} a^{D-2} - \frac{1}{2} Z m_0^2 \varphi^2 a^D - \frac{1}{4!} Z^2 \lambda_0 \varphi^4 a^D. \quad (21)$$

We now make the usual decomposition of bare parameters into renormalized parameters and counterterms,

$$Z \equiv 1 + \delta Z, \quad Z m_0^2 \equiv 0 + \delta m^2, \quad Z^2 \lambda_0 \equiv \lambda + \delta \lambda. \quad (22)$$

Note that we define the renormalized mass to be zero. Of course this can only be enforced initially; the point of the calculation is to see what happens later under the impact of inflationary particle production!

The final expression of the Lagrangian distinguishes the free part (1st term), the interaction (2nd term) and the counter-interactions (last 3 terms),

$$\mathcal{L} = -\frac{1}{2} \partial_\mu \varphi \partial_\nu \varphi \eta^{\mu\nu} a^{D-2} - \frac{1}{4!} \lambda \varphi^4 a^D - \frac{1}{2} \delta Z \partial_\mu \varphi \partial_\nu \varphi \eta^{\mu\nu} a^{D-2} - \frac{1}{2} \delta m^2 \varphi^2 a^D - \frac{1}{4!} \delta \lambda \varphi^4 a^D. \quad (23)$$

The propagator derives from the free part and obeys the equation,

$$\text{Propagator} \quad \Longrightarrow \quad \eta^{\mu\nu} \partial_\mu \left(a^{D-2} \partial_\nu i\Delta(x; x') \right) = i\delta^D(x - x') . \quad (24)$$

The position-space interaction vertex depends upon the four points x_{1-4}^μ ,

$$\text{Interaction} \quad \Longrightarrow \quad -i\lambda a_1^D \delta^D(x_1 - x_2) \delta^D(x_2 - x_3) \delta^D(x_3 - x_4) . \quad (25)$$

The field strength renormalization depends upon two points x^μ and x'^μ ,

$$\text{Field Strength Renormalization} \quad \Longrightarrow \quad i\delta Z \eta^{\mu\nu} \partial_\mu \left(a^{D-2} \partial_\nu \delta^D(x - x') \right) . \quad (26)$$

The mass renormalization also depends upon x^μ and x'^μ ,

$$\text{Mass Renormalization} \quad \Longrightarrow \quad -i\delta m^2 a^D \delta^D(x - x') . \quad (27)$$

Of course the coupling constant renormalization is the same as (25) with λ replaced by $\delta\lambda$,

$$\text{Coupling Constant Renorm.} \quad \Longrightarrow \quad -i\delta\lambda a_1^D \delta^D(x_1 - x_2) \delta^D(x_2 - x_3) \delta^D(x_3 - x_4) . \quad (28)$$

It remains to solve (24) for the propagator. One might think to look for a function of the de Sitter invariant geodesic length $\ell(x; x')$ but there is no such solution. The reason is that the free theory possesses no normalizable, de Sitter invariant wave functions, and hence no de Sitter invariant states [86]. By allowing $i\Delta(x; x')$ to depend additionally upon the scale factors $a \equiv a(\eta)$ and $a' \equiv a(\eta')$ one can find a solution which has the homogeneity and isotropy of cosmology [2]. It is expressed in terms of a length function we call $y_{++}(x; x')$,

$$y_{++}(x; x') \equiv aa' H^2 \left[\|\vec{x} - \vec{x}'\|^2 - (|\eta - \eta'| - i\delta)^2 \right] \equiv aa' H^2 \Delta x_{++}^2(x; x') , \quad (29)$$

When the infinitesimal parameter δ vanishes this has a simple relation to the geodesic length, $y = 4 \sin^2(H\ell/2)$. One can see from (13-16) that the four propagators must all depend the same way on a , a' and length functions $y_{\pm\pm}$ defined as follows,

$$y_{+-}(x; x') \equiv aa' H^2 \left[\|\vec{x} - \vec{x}'\|^2 - (\eta - \eta' + i\delta)^2 \right] \equiv aa' H^2 \Delta x_{+-}^2(x; x') , \quad (30)$$

$$y_{-+}(x; x') \equiv aa' H^2 \left[\|\vec{x} - \vec{x}'\|^2 - (\eta - \eta' - i\delta)^2 \right] \equiv aa' H^2 \Delta x_{-+}^2(x; x') , \quad (31)$$

$$y_{--}(x; x') \equiv aa' H^2 \left[\|\vec{x} - \vec{x}'\|^2 - (|\eta - \eta'| + i\delta)^2 \right] \equiv aa' H^2 \Delta x_{--}^2(x; x') . \quad (32)$$

Because of this close relationship between the four propagators we shall often drop the $\pm$ subscripts and treat expressions for general $y = y(x; x')$ and $\Delta x^2 = \Delta x^2(x; x')$.

To facilitate dimensional regularization we express the dimension of spacetime in terms of its deviation from four, $D = 4 - \epsilon$. It is also convenient to break the propagator up into three terms,

$$i\Delta(x; x') = A(x; x') + B(x; x') + C(x; x') , \quad (33)$$

$$A(x; x') \equiv \frac{\Gamma(1-\frac{\epsilon}{2}) (aa')^{-1+\frac{\epsilon}{2}}}{4\pi^{2-\frac{\epsilon}{2}} \Delta x^{2-\epsilon}} , \quad (34)$$

$$B(x; x') \equiv \frac{H^{2-\epsilon}}{(4\pi)^{2-\frac{\epsilon}{2}}} \left\{ -\frac{2\Gamma(3-\frac{\epsilon}{2})}{\epsilon} \left(\frac{y}{4}\right)^{\frac{\epsilon}{2}} + \pi \cot\left(\frac{\pi\epsilon}{2}\right) \frac{\Gamma(3-\epsilon)}{\Gamma(2-\frac{\epsilon}{2})} + \frac{\Gamma(3-\epsilon)}{\Gamma(2-\frac{\epsilon}{2})} \ln(aa') \right\} , \quad (35)$$

$$C(x; x') \equiv \frac{H^{2-\epsilon}}{(4\pi)^{2-\frac{\epsilon}{2}}} \sum_{n=1}^{\infty} \left[\frac{1}{n} \frac{\Gamma(3-\epsilon+n)}{\Gamma(2-\frac{\epsilon}{2}+n)} \left(\frac{y}{4}\right)^n - \frac{1}{n+\frac{\epsilon}{2}} \frac{\Gamma(3-\frac{\epsilon}{2}+n)}{\Gamma(2+n)} \left(\frac{y}{4}\right)^{n+\frac{\epsilon}{2}} \right] . \quad (36)$$

Powers of $A(x; x')$ are the ultimate source of ultraviolet divergences. The term $B(x; x')$ is not ultraviolet divergent by itself,

$$\lim_{\epsilon \rightarrow 0} B(x; x') = \left(\frac{H}{4\pi}\right)^2 \left\{ -2 \ln(H^2 \Delta x^2 / 4) - 1 \right\} = -\frac{H^2}{8\pi^2} \ln\left(\frac{\sqrt{e}}{4} H^2 \Delta x^2\right). \quad (37)$$

However, we must retain $\epsilon \neq 0$ whenever $B(x; x')$ multiplies two or more powers of $A(x; x')$. The term $C(x; x')$ actually vanishes for $\epsilon = 0$. We only need it when multiplying by more than two powers of $A(x; x')$, which can not happen for the maximum number of propagators — $(i\Delta(x; x'))^3$ — we shall encounter in this computation.

III. ONE LOOP COUNTERTERMS

Neither the non-flat geometry, nor the Schwinger-Keldysh formalism, nor even the use of position space rather than momentum space, alters the basic structure of perturbative quantum field theory. In particular, the diagrams all look the same as in flat momentum space, and working at two loops requires knowledge of the one loop counterterms. The purpose of this section is to compute the order λ contribution to δm^2 and the order λ^2 contribution to $\delta\lambda$. As in flat space there is no one loop contribution to δZ .

Fig. 1 depicts the one loop contribution to the scalar self-mass-squared. We can extract δm^2 at order λ by focusing on the $++$ term,

$$-iM_{1++}^2(x; x') = -i \left[\frac{\lambda}{2} i\Delta_{++}(x; x) + \delta m^2 \right] a^D \delta^D(x-x') + O(\lambda^2) . \quad (38)$$

Although the coincidence limit of the massless propagator vanishes when using dimensional regularization in flat space, the de Sitter curvature of our background gives a nonzero result




FIG. 1: Generic one loop self-mass-squared.

that can be read off from (33),

$$i\Delta(x'; x') = \lim_{x \rightarrow x'} i\Delta(x; x') = \frac{H^{2-\epsilon}}{(4\pi)^{2-\frac{\epsilon}{2}}} \frac{\Gamma(3-\epsilon)}{\Gamma(2-\frac{\epsilon}{2})} \left\{ 2 \ln(a) + \pi \cot\left(\frac{\pi\epsilon}{2}\right) \right\}. \quad (39)$$

Because of the finite, time-dependent term we cannot make the one loop self-mass-squared vanish for all time. Our renormalization condition is that it should be zero at $t = 0$, which implies,

$$\delta m^2 = -\frac{\lambda H^{2-\epsilon}}{2^{5-\epsilon} \pi^{2-\frac{\epsilon}{2}}} \frac{\Gamma(3-\epsilon)}{\Gamma(2-\frac{\epsilon}{2})} \pi \cot\left(\frac{\pi\epsilon}{2}\right) + O(\lambda^2). \quad (40)$$

The renormalized $++$ self-mass-squared is therefore,

$$M_{1++}^2(x; x') = \frac{\lambda H^2}{8\pi^2} a^4 \ln(a) \delta^4(x - x') + O(\lambda^2). \quad (41)$$

The $--$ case is minus the complex conjugate,

$$M_{1--}^2(x; x') = -\frac{\lambda H^2}{8\pi^2} a^4 \ln(a) \delta^4(x - x') + O(\lambda^2). \quad (42)$$

The $+-$ and $-+$ cases vanish at this order because there are no mixed interactions,

$$M_{1+-}^2(x; x') = -\left(M_{1-+}^2(x; x')\right)^* = 0 + O(\lambda^2). \quad (43)$$




FIG. 2: Generic tree and one loop vertex function.

We can extract the order λ^2 contribution to $\delta\lambda$ from the one loop corrections to the $++++$ 4-point 1PI function. Fig. 2 shows the diagram topology. A simple application of the

Feynman rules gives,

$$\begin{aligned}
-iV_+(x_1, x_2, x_3, x_4) = & -i(\lambda + \delta\lambda) a_1^D \delta^D(x_1 - x_2) \delta^D(x_1 - x_3) \delta^D(x_1 - x_4) - \frac{1}{2} \lambda^2 \left\{ \right. \\
& a_1^D a_4^D \left[i\Delta_{++}(x_1; x_4) \right]^2 \delta^D(x_1 - x_2) \delta^D(x_3 - x_4) + a_1^D a_3^D \left[i\Delta_{++}(x_1; x_3) \right]^2 \delta^D(x_1 - x_4) \delta^D(x_2 - x_3) \\
& \left. + a_1^D a_2^D \left[i\Delta_{++}(x_1; x_2) \right]^2 \delta^D(x_1 - x_3) \delta^D(x_4 - x_2) \right\} + O(\lambda^3) . \quad (44)
\end{aligned}$$

In Appendix B we show that the square of a propagator can be expressed as a local term which diverges for $\epsilon \rightarrow 0$, plus a nonlocal term which is integrable for $\epsilon = 0$. For the $++$ propagator the result is equation (95) which we reproduce here,

$$\begin{aligned}
(i\Delta_{++}(x; x'))^2 \longrightarrow & -\frac{i\mu^{-\epsilon}}{2^3\pi^{2-\frac{\epsilon}{2}}} \frac{\Gamma(1-\frac{\epsilon}{2})}{(1-\epsilon)\epsilon} a^{-4+2\epsilon} \delta^D(x-x') - \frac{(aa')^{-2}}{2^6\pi^4} \partial^2 \left(\frac{\ln(\mu^2 \Delta x_{++}^2)}{\Delta x_{++}^2} \right) \\
& - \frac{H^2}{2^4\pi^4} \frac{\ln\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{++}^2\right)}{aa' \Delta x_{++}^2} + \frac{H^4}{2^6\pi^4} \ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{++}^2\right). \quad (45)
\end{aligned}$$

To fix $\delta\lambda$ we are interested only in the local, divergent term. There is such a contribution from the s -channel — $(i\Delta_{++}(x_1; x_4))^2$ — the t -channel — $(i\Delta_{++}(x_1; x_3))^2$ — and the u -channel — $(i\Delta_{++}(x_1; x_2))^2$ — graphs. They each make the same contribution, so the divergent part of the vertex is,

$$\begin{aligned}
-iV_+(x_1, x_2, x_3, x_4) = & -i(\lambda + \delta\lambda) a_1^{4-\epsilon} \delta^D(x_1 - x_2) \delta^D(x_1 - x_3) \delta^D(x_1 - x_4) \\
& + \frac{i3\lambda^2 \mu^{-\epsilon}}{2^4\pi^{2-\frac{\epsilon}{2}}} \frac{\Gamma(1-\frac{\epsilon}{2})}{(1-\epsilon)\epsilon} a_1^4 \delta^D(x_1 - x_2) \delta^D(x_1 - x_3) \delta^D(x_1 - x_4) \\
& + \lambda^2 \times (\text{Nonlocal Finite}) + O(\lambda^3) . \quad (46)
\end{aligned}$$

The simplest renormalization condition is to absorb the entire local, constant part at order λ^2 ,

$$\delta\lambda = \frac{3\lambda^2 \mu^{-\epsilon}}{16\pi^{2-\frac{\epsilon}{2}}} \frac{\Gamma(1-\frac{\epsilon}{2})}{\epsilon(1-\epsilon)} . \quad (47)$$

Note that while the divergence is proportional to the fourth power of the scale factor, the counterterm is proportional to the D -th power. This means that expanding $a_1^{4-\epsilon} = a_1^4(1 - \epsilon \ln(a_1) + O(\epsilon^2))$, gives an additional, finite local term. The full renormalized $++++$ 1PI 4-point vertex is,

$$\begin{aligned}
-iV_R(x_1, x_2, x_3, x_4) = & -i \left(\lambda - \frac{3\lambda^2}{2^4\pi^2} \ln(a_1) \right) a_1^4 \delta^4(x_1 - x_2) \delta^4(x_1 - x_3) \delta^4(x_1 - x_4) + \frac{\lambda^2}{2^7\pi^4} \left\{ \right. \\
& (a_1 a_4)^2 \partial_1^2 \left[\frac{\ln(\mu^2 \Delta x_{14}^2)}{\Delta x_{14}^2} \right] + 4H^2 (a_1 a_4)^3 \frac{\ln\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{14}^2\right)}{\Delta x_{14}^2} - H^4 (a_1 a_4)^4 \ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{14}^2\right) \left. \right\} \\
& \times \delta^4(x_1 - x_2) \delta^4(x_3 - x_4) + \lambda^2 (t\text{-channel}) + \lambda^2 (u\text{-channel}) + O(\lambda^3) . \quad (48)
\end{aligned}$$

IV. THE SUNSET DIAGRAM

The first two loop contribution to the scalar self-mass-squared is known as the sunset (ss) diagram. Fig. 3 depicts its topology. A simple application of the Feynman rules gives,

$$-iM_{ss\pm\pm}^2 = \frac{1}{3!}(\mp i\lambda)(\mp i\lambda)(aa')^{4-\epsilon} \left(i\Delta_{\pm\pm}(x; x') \right)^3. \quad (49)$$




FIG. 3: Generic Sunset Diagram.

In Appendix C we show that the cube of a propagator can be expressed as a local term which diverges for $\epsilon \rightarrow 0$, plus a nonlocal term which is integrable for $\epsilon = 0$. For the $++$ propagator the result is equation (108) which we reproduce here,

$$\begin{aligned} (i\Delta_{++}(x; x'))^3 \rightarrow & -\frac{i\mu^{-2\epsilon}\Gamma^2(1-\frac{\epsilon}{2})}{2^9\pi^{4-\epsilon}} \frac{(aa')^{-3+\frac{3}{2}\epsilon}\partial^2\delta^D(x-x')}{(1-\frac{3}{2}\epsilon)(1-\epsilon)(1-\frac{3}{4}\epsilon)\epsilon} + \frac{i3H^{2-\epsilon}\mu^{-\epsilon}a^{-4+2\epsilon}}{2^{6-\epsilon}\pi^{4-\epsilon}} \left\{ \right. \\ & \frac{(1-\frac{\epsilon}{2})(1-\frac{\epsilon}{4})\Gamma^2(1-\frac{\epsilon}{2})}{(1-\frac{3}{2}\epsilon)\epsilon^2} \left(\frac{Ha}{2\mu} \right)^\epsilon - \frac{\Gamma(1-\epsilon)}{\epsilon} \left[\pi \cot\left(\frac{\pi\epsilon}{2}\right) + 2\ln(a) \right] \left. \right\} \delta^D(x-x') \\ & - \frac{(aa')^{-3}}{2^{11}\pi^6} \partial^4 \left[\frac{\ln(\mu^2\Delta_{++}^2)}{\Delta_{++}^2} \right] + \frac{3H^2(aa')^{-2}}{2^8\pi^6} \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2\Delta_{++}^2)}{\Delta_{++}^2} + \frac{1}{4} \frac{\ln^2(\mu^2\Delta_{++}^2)}{\Delta_{++}^2} \right] \\ & + \frac{3H^4\ln^2\left(\frac{\sqrt{e}}{4}H^2\Delta_{++}^2\right)}{2^8\pi^6 aa'\Delta_{++}^2} - \frac{H^6}{2^9\pi^6} \ln^3\left(\frac{\sqrt{e}}{4}H^2\Delta_{++}^2\right). \quad (50) \end{aligned}$$

When this is multiplied by $(aa')^{4-\epsilon}$ a few simple rearrangements allow us to recognize the spacetime dependent part of the the first term as essentially a field strength renormalization,

$$\begin{aligned} (aa')^{1+\frac{\epsilon}{2}}\partial^2\delta^D(x-x') &= a^{1+\frac{\epsilon}{2}}\partial^2\left(a^{1+\frac{\epsilon}{2}}\delta^D(x-x')\right) \\ &= a^{2+\epsilon} \left\{ \partial^2 - (2+\epsilon)Ha\partial_0 - \left(1+\frac{\epsilon}{2}\right)\left(2+\frac{\epsilon}{2}\right)H^2a^2 \right\} \delta^D(x-x'). \quad (51) \end{aligned}$$

Doing the same thing for the second term reveals what is essentially a mass renormalization plus some finite local terms *and* an overlapping divergence,

$$\frac{i3H^2a^4}{2^6\pi^4} \left(\frac{2\pi}{H\mu} \right)^\epsilon \left\{ \frac{(1-\frac{\epsilon}{4})(1-\frac{\epsilon}{2})\Gamma^2(1-\frac{\epsilon}{2})}{\epsilon^2(1-\frac{3}{2}\epsilon)} \left(\frac{Ha}{2\mu} \right)^\epsilon - \frac{\Gamma(1-\epsilon)}{\epsilon} \left[\pi \cot\left(\frac{\pi\epsilon}{2}\right) + 2\ln(a) \right] \right\} \delta^D(x-x')$$

$$\begin{aligned} \longrightarrow \frac{i3H^2a^4}{2^6\pi^4} \left(\frac{2\pi}{H\mu} \right)^\epsilon & \left\{ \frac{(1-\frac{\epsilon}{4})(1-\frac{\epsilon}{2})\Gamma^2(1-\frac{\epsilon}{2})}{\epsilon^2(1-\frac{3}{2}\epsilon)} \left(\frac{H}{2\mu} \right)^\epsilon - \frac{\Gamma(1-\epsilon)}{\epsilon} \pi \cot\left(\frac{\pi\epsilon}{2}\right) \right. \\ & \left. - \frac{1}{\epsilon} \ln(a) + \frac{1}{2} \ln^2(a) + \left[\frac{3}{4} - \gamma + \ln\left(\frac{H}{2\mu}\right) \right] \ln(a) \right\} \delta^D(x-x'). \end{aligned} \quad (52)$$

In displaying the final result we exhibit the field strength and mass renormalizations, as well as the overlapping divergence and the finite, local terms. Our result for the ++ case is,

$$\begin{aligned} M_{ss++}^2 \longrightarrow & -\frac{\lambda^2}{2^{10}3\pi^4} \left(\frac{\pi}{\mu^2} \right)^\epsilon \frac{\Gamma^2(1-\frac{\epsilon}{2})}{(1-\frac{3}{4}\epsilon)(1-\frac{3}{2}\epsilon)(1-\epsilon)\epsilon} a^2 [\partial^2 - 2Ha\partial_0 - 2H^2a^2] \delta^D(x-x') \\ & - \frac{\lambda^2}{2^{10}3\pi^4} a^2 \left\{ \ln(a)\partial^2 - (2\ln(a)+1)Ha\partial_0 - \left(2\ln(a)+\frac{3}{2}\right)H^2a^2 \right\} \delta^4(x-x') \\ & + \frac{\lambda^2H^2}{2^7\pi^4} \left(\frac{2\pi}{H\mu} \right)^\epsilon \left\{ \frac{(1-\frac{\epsilon}{4})(1-\frac{\epsilon}{2})\Gamma^2(1-\frac{\epsilon}{2})}{\epsilon^2(1-\frac{3}{2}\epsilon)} \left(\frac{H}{2\mu} \right)^\epsilon - \frac{\Gamma(1-\epsilon)}{\epsilon} \pi \cot\left(\frac{\pi\epsilon}{2}\right) \right\} a^4 \delta^D(x-x') \\ & - \frac{\lambda^2H^2}{2^7\pi^4} \left(\frac{2\pi}{H\mu} \right)^\epsilon \frac{\ln(a)}{\epsilon} a^4 \delta^D(x-x') + \frac{\lambda^2H^2}{2^7\pi^4} \left\{ \frac{1}{2} \ln^2(a) + \left[\frac{3}{4} - \gamma + \ln\left(\frac{H}{2\mu}\right) \right] \ln(a) \right\} a^4 \delta^4(x-x') \\ & + \frac{i\lambda^2aa'}{2^9\pi^6} \left\{ \frac{1}{24} \partial^4 \left[\frac{\ln(\mu^2\Delta x_{++}^2)}{\Delta x_{++}^2} \right] - H^2aa' \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2\Delta x_{++}^2)}{\Delta x_{++}^2} + \frac{\ln^2(\mu^2\Delta x_{++}^2)}{4\Delta x_{++}^2} \right] \right. \\ & \quad \left. - H^4(aa')^2 \frac{\ln^2\left(\frac{\sqrt{e}}{4}H^2\Delta x_{++}^2\right)}{\Delta x_{++}^2} + \frac{H^6}{6} (aa')^3 \ln^3\left(\frac{\sqrt{e}}{4}H^2\Delta x_{++}^2\right) \right\}. \end{aligned} \quad (53)$$

The -- case is minus the complex conjugate,

$$\begin{aligned} M_{ss--}^2 \longrightarrow & \frac{\lambda^2}{2^{10}3\pi^4} \left(\frac{\pi}{\mu^2} \right)^\epsilon \frac{\Gamma^2(1-\frac{\epsilon}{2})}{(1-\frac{3}{4}\epsilon)(1-\frac{3}{2}\epsilon)(1-\epsilon)\epsilon} a^2 [\partial^2 - 2Ha\partial_0 - 2H^2a^2] \delta^D(x-x') \\ & + \frac{\lambda^2}{2^{10}3\pi^4} a^2 \left\{ \ln(a)\partial^2 - (2\ln(a)+1)Ha\partial_0 - \left(2\ln(a)+\frac{3}{2}\right)H^2a^2 \right\} \delta^4(x-x') \\ & - \frac{\lambda^2H^2}{2^7\pi^4} \left(\frac{2\pi}{H\mu} \right)^\epsilon \left\{ \frac{(1-\frac{\epsilon}{4})(1-\frac{\epsilon}{2})\Gamma^2(1-\frac{\epsilon}{2})}{\epsilon^2(1-\frac{3}{2}\epsilon)} \left(\frac{H}{2\mu} \right)^\epsilon - \frac{\Gamma(1-\epsilon)}{\epsilon} \pi \cot\left(\frac{\pi\epsilon}{2}\right) \right\} a^4 \delta^D(x-x') \\ & + \frac{\lambda^2H^2}{2^7\pi^4} \left(\frac{2\pi}{H\mu} \right)^\epsilon \frac{\ln(a)}{\epsilon} a^4 \delta^D(x-x') - \frac{\lambda^2H^2}{2^7\pi^4} \left\{ \frac{1}{2} \ln^2(a) + \left[\frac{3}{4} - \gamma + \ln\left(\frac{H}{2\mu}\right) \right] \ln(a) \right\} a^4 \delta^4(x-x') \\ & + \frac{i\lambda^2aa'}{2^9\pi^6} \left\{ \frac{1}{24} \partial^4 \left[\frac{\ln(\mu^2\Delta x_{--}^2)}{\Delta x_{--}^2} \right] - H^2aa' \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2\Delta x_{--}^2)}{\Delta x_{--}^2} + \frac{\ln^2(\mu^2\Delta x_{--}^2)}{4\Delta x_{--}^2} \right] \right. \\ & \quad \left. - H^4(aa')^2 \frac{\ln^2\left(\frac{\sqrt{e}}{4}H^2\Delta x_{--}^2\right)}{\Delta x_{--}^2} + \frac{H^6}{6} (aa')^3 \ln^3\left(\frac{\sqrt{e}}{4}H^2\Delta x_{--}^2\right) \right\}. \end{aligned} \quad (54)$$

The mixed polarities lack the local terms and differ by a sign,

$$\begin{aligned} M_{ss+-}^2 = & \frac{i\lambda^2}{6} (aa')^{4-\epsilon} [i\Delta_{+-}(x; x')]^3 = -i \frac{\lambda^2aa'}{2^9\pi^6} \left\{ \frac{1}{24} \partial^4 \left[\frac{\ln(\mu^2\Delta x_{+-}^2)}{\Delta x_{+-}^2} \right] \right. \\ & \left. - H^2aa' \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2\Delta x_{+-}^2)}{\Delta x_{+-}^2} + \frac{1}{4} \frac{\ln^2(\mu^2\Delta x_{+-}^2)}{\Delta x_{+-}^2} \right] - H^4(aa')^2 \frac{\ln^2\left(\frac{\sqrt{e}}{4}H^2\Delta x_{+-}^2\right)}{\Delta x_{+-}^2} \right\} \end{aligned}$$

$$+\frac{H^6}{6}(aa')^3 \ln^3 \left(\frac{\sqrt{e}}{4} H^2 \Delta x_{+-}^2 \right) \Big\} , \quad (55)$$

$$\begin{aligned} M_{\text{ss}-+}^2 &= \frac{i\lambda^2}{6}(aa')^{4-\epsilon} [i\Delta_{-+}(x; x')]^3 = -i \frac{\lambda^2 aa'}{2^9 \pi^6} \left\{ \frac{1}{24} \partial^4 \left[\frac{\ln(\mu^2 \Delta x_{-+}^2)}{\Delta x_{-+}^2} \right] \right. \\ &\quad \left. - H^2 aa' \partial^2 \left[\ln \left(\frac{H e^{\frac{3}{4}}}{2\mu} \right) \frac{\ln(\mu^2 \Delta x_{-+}^2)}{\Delta x_{-+}^2} + \frac{1}{4} \frac{\ln^2(\mu^2 \Delta x_{-+}^2)}{\Delta x_{-+}^2} \right] - H^4 (aa')^2 \frac{\ln^2 \left(\frac{\sqrt{e}}{4} H^2 \Delta x_{-+}^2 \right)}{\Delta x_{-+}^2} \right. \\ &\quad \left. + \frac{H^6}{6} (aa')^3 \ln^3 \left(\frac{\sqrt{e}}{4} H^2 \Delta x_{-+}^2 \right) \right\} . \quad (56) \end{aligned}$$

V. THE SNOWMAN DIAGRAM

The primitive two loop graph is known as the snowman (sm) diagram. Unlike the sunset diagram, it contributes only to M_{++}^2 and M_{--}^2 . Its generic topology is depicted in Fig. 4. The right hand diagram depicts the 1-loop mass counterterm which naturally combines with it.




FIG. 4: Generic Snowman Diagram with mass counterterm.

Because both external points of this diagram must have the same polarity we denote the contributions with a single subscript sign, $M_{\text{sm}+}^2$ and $M_{\text{sm}-}^2$. The internal vertices are summed over both polarities, as always in the Schwinger-Keldysh formalism. A simple application of the Feynman rules gives,

$$\begin{aligned} -iM_{\text{sm}+}^2 &= \frac{-i\lambda a^D}{2} \delta^D(x-x') \int d^D x'' a''^D \left\{ [i\Delta_{++}(x; x'')]^2 - [i\Delta_{+-}(x; x'')]^2 \right\} \\ &\quad \times \left\{ \frac{(-i\lambda)}{2} i\Delta(x''; x'') - i\delta m^2 \right\} . \quad (57) \end{aligned}$$

Since both diagrams of Fig. 4 have the same lower loop, they possess a common factor in the first curly bracket. The first term within the final curly bracket derives from the left hand diagram, while the second term comes from the right hand diagram.

The coincident propagator and the mass counter term are calculated in (39) and (40) respectively. Once they are inserted into the above equation, one finds

$$-iM_{\text{sm}+}^2 = -\frac{\lambda^2 H^{2-\epsilon} a^{4-\epsilon}}{2^{5-\epsilon} \pi^{2-\frac{\epsilon}{2}}} \frac{\Gamma(3-\epsilon)}{\Gamma(2-\frac{\epsilon}{2})} \delta^D(x-x') \int d^4 x'' a''^{4-\epsilon} \ln(a'') \left\{ [i\Delta_{++}(x; x'')]^2 - [i\Delta_{+-}(x; x'')]^2 \right\}. \quad (58)$$

The squares of the various propagators are computed in Appendix B and the results we need are (95) and (96)

$$\begin{aligned} -iM_{\text{sm}+}^2 &= \frac{i\lambda^2 H^2 a^4}{2^7 \pi^4} \left(\frac{2\pi}{H\mu} \right)^\epsilon \frac{\Gamma(1-\epsilon)}{\epsilon} \ln(a) \delta^D(x-x') \\ &\quad + \frac{\lambda^2 H^2 a^2}{2^{10} \pi^6} \delta^4(x-x') (-\partial_0^2) \int d^4 x'' a''^2 \ln(a'') \left[\frac{\ln(\mu^2 \Delta x_{++}^2)}{\Delta x_{++}^2} - \frac{\ln(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} \right] \\ &\quad + \frac{\lambda^2 H^4 a^3}{2^8 \pi^6} \delta^4(x-x') \int d^4 x'' a''^3 \ln(a'') \left[\frac{\ln(\frac{H^2 \sqrt{e}}{4} \Delta x_{++}^2)}{\Delta x_{++}^2} - \frac{\ln(\frac{H^2 \sqrt{e}}{4} \Delta x_{+-}^2)}{\Delta x_{+-}^2} \right] \\ &\quad - \frac{\lambda^2 H^6 a^4}{2^{10} \pi^6} \delta^4(x-x') \int d^4 x'' a''^4 \ln(a'') \left[\ln^2\left(\frac{H^2 \sqrt{e}}{4} \Delta x_{++}^2\right) - \ln^2\left(\frac{H^2 \sqrt{e}}{4} \Delta x_{+-}^2\right) \right], \quad (59) \\ &\longrightarrow \frac{i\lambda^2 H^2}{2^7 \pi^4} \left(\frac{2\pi}{H\mu} \right)^\epsilon \left\{ \frac{\ln(a)}{\epsilon} + \gamma \ln(a) \right\} a^4 \delta^D(x-x') \\ &\quad + \frac{\lambda^2 H^2 a^2}{2^{10} \pi^6} \delta^4(x-x') \left\{ I_1 + 4H^2 a I_2 - H^4 a^2 I_3 \right\}. \quad (60) \end{aligned}$$

The integrals I_1 , I_2 and I_3 are evaluated in Appendix D and the results are expressions (119), (125), and (129). Substituting these and adding everything up gives,

$$\begin{aligned} M_{\text{sm}+}^2 &= -\frac{\lambda^2 H^2}{2^7 \pi^4} \left(\frac{2\pi}{H\mu} \right)^\epsilon \frac{\ln(a)}{\epsilon} a^4 \delta^D(x-x') + \frac{\lambda^2 H^2}{2^7 \pi^4} a^4 \delta^4(x-x') \left\{ -\frac{4}{9} \ln^3(a) \right. \\ &\quad + \frac{13}{18} \ln^2(a) + \left[\ln\left(\frac{H}{2\mu}\right) + \frac{8}{3} - \gamma - \frac{2}{9} \pi^2 \right] \ln(a) - \frac{238}{81} + \frac{13}{54} \pi^2 + \frac{4}{3} \zeta(3) + \frac{a^{-3}}{81} \\ &\quad \left. - \sum_{n=1}^{\infty} \frac{n+5}{(n+1)^3} a^{-(n+1)} + 4 \sum_{n=1}^{\infty} \frac{a^{-(n+2)}}{(n+2)^3} + 4 \sum_{n=1}^{\infty} \frac{a^{-(n+3)}}{n(n+3)^3} \right\}. \quad (61) \end{aligned}$$

Note that the overlapping divergence from the first term is the same as the one from the sunset diagram. The -- case is minus (61).

VI. THE COUNTER DIAGRAMS

We have already encountered the mass counterterm in Section III. At the order we are working there is also a one loop diagram with λ replaced by $\delta\lambda$, and there is a field strength




FIG. 5: Generic Counter Diagrams. The one on the left is the same as Fig. 1 but with $\delta\lambda$ in place of λ . The one on the right is the field strength renormalization.

renormalization. The generic diagram topologies are depicted in Fig. 5. Note that neither diagram contributes to the mixed polarities.

The contribution of the one loop counter vertex diagram to the $++$ polarity is,

$$-iM_{\text{cv}+}^2 = -i\frac{\delta\lambda}{2}a^D i\Delta(x'; x')\delta^D(x-x'). \quad (62)$$

Of course we can read off the answer from Section III,

$$M_{\text{cv}+}^2 = \frac{3H^2\lambda^2a^{4-\epsilon}}{2^8\pi^4}\left(\frac{2\pi}{\mu H}\right)^\epsilon \frac{\Gamma(1-\epsilon)}{\epsilon} \left\{2\ln(a) + \pi \cot\left(\frac{\pi\epsilon}{2}\right)\right\} \delta^D(x-x'), \quad (63)$$

$$\longrightarrow \frac{H^2\lambda^2}{2^7\pi^4}\left(\frac{2\pi}{\mu H}\right)^\epsilon \left\{\frac{3}{2}\frac{\Gamma(1-\epsilon)}{\epsilon}\pi \cot\left(\frac{\pi\epsilon}{2}\right) - \frac{3}{2}\ln^2(a)\right\} a^4\delta^D(x-x'). \quad (64)$$

The $--$ case is minus this.

The contribution of field strength renormalization to the $++$ polarity is,

$$M_{\delta Z+}^2 = -\delta Z \eta^{\mu\nu} \partial_\mu (a^{D-2} \partial_\nu \delta^D(x-x')) = -\delta Z a^{2-\epsilon} (\partial^2 - (2-\epsilon)Ha\partial_0) \delta^D(x-x'). \quad (65)$$

The only primitive graph with a divergence proportional to $a^2(\partial^2 - 2Ha\partial_0)\delta^D(x-x')$ is the sunset diagram. Comparison with our result (53) for it implies,

$$\delta Z = -\frac{\lambda^2}{2^{10}3\pi^4}\left(\frac{\pi}{\mu^2}\right)^\epsilon \frac{\Gamma^2(1-\frac{\epsilon}{2})}{(1-\frac{3}{4}\epsilon)(1-\frac{3}{2}\epsilon)(1-\epsilon)\epsilon}. \quad (66)$$

Although some finite, local terms result from expanding $a^{-\epsilon} = 1 - \epsilon \ln(a) + O(\epsilon^2)$ there are no overlapping divergences from the field strength renormalization at two loop order,

$$\begin{aligned} M_{\delta Z+}^2 &= \frac{\lambda^2}{2^{10}3\pi^4}\left(\frac{\pi}{\mu^2}\right)^\epsilon \frac{\Gamma^2(1-\frac{\epsilon}{2})}{(1-\frac{3}{4}\epsilon)(1-\frac{3}{2}\epsilon)(1-\epsilon)\epsilon} a^2 (\partial^2 - 2Ha\partial_0) \delta^D(x-x') \\ &\quad + \frac{\lambda^2}{2^{10}3\pi^4} a^2 \left[-\ln(a)\partial^2 + (2\ln(a)+1)Ha\partial_0 \right] \delta^4(x-x'). \end{aligned} \quad (67)$$

After field strength renormalization, all remaining divergences reside on $a^4\delta^D(x-x')$. Such contributions comes from the sunset diagram (53) and the counter vertex (64). Comparison with these results suggests that the simplest choice for the two loop mass counterterm is,

$$\delta m^2|_{\lambda^2} = -\frac{\lambda^2 H^2}{2^7 \pi^4} \left(\frac{2\pi}{H\mu} \right)^\epsilon \left\{ \left(\frac{H}{2\mu} \right)^\epsilon \frac{\Gamma^2(1-\frac{\epsilon}{2})}{12(1-\frac{3}{2}\epsilon)(1-\frac{3}{4}\epsilon)(1-\epsilon)\epsilon} + \left(\frac{H}{2\mu} \right)^\epsilon \frac{(1-\frac{\epsilon}{4})(1-\frac{\epsilon}{2})\Gamma^2(1-\frac{\epsilon}{2})}{(1-\frac{3}{2}\epsilon)\epsilon^2} \right. \\ \left. + \frac{\Gamma(1-\epsilon)}{2\epsilon} \pi \cot\left(\frac{\pi\epsilon}{2}\right) + \frac{1}{16} - \frac{238}{81} + \frac{13}{54}\pi^2 + \frac{4}{3}\zeta(3) \right\}. \quad (68)$$

Of course the mass counterterm is $M_{\delta m+}^2 = \delta m^2 a^{4-\epsilon} \delta^D(x-x')$. Using the expansion $a^{-\epsilon} = 1 - \epsilon \ln(a) + \frac{1}{2}\epsilon^2 \ln^2(a) + O(\epsilon^3)$ we see that the two loop contribution is,

$$M_{\delta m+}^2 \Big|_{\lambda^2} \longrightarrow -\frac{\lambda^2 H^2}{2^7 \pi^4} \left(\frac{2\pi}{H\mu} \right)^\epsilon \left\{ \left(\frac{H}{2\mu} \right)^\epsilon \frac{\Gamma^2(1-\frac{\epsilon}{2})}{12(1-\frac{3}{2}\epsilon)(1-\frac{3}{4}\epsilon)(1-\epsilon)\epsilon} + \left(\frac{H}{2\mu} \right)^\epsilon \frac{(1-\frac{\epsilon}{4})(1-\frac{\epsilon}{2})\Gamma^2(1-\frac{\epsilon}{2})}{(1-\frac{3}{2}\epsilon)\epsilon^2} \right. \\ \left. + \frac{\Gamma(1-\epsilon)}{2\epsilon} \pi \cot\left(\frac{\pi\epsilon}{2}\right) \right\} a^4 \delta^D(x-x') + \frac{\lambda^2 H^2}{2^6 \pi^4} \left(\frac{2\pi}{H\mu} \right)^\epsilon \frac{\ln(a)}{\epsilon} a^4 \delta^D(x-x') \\ - \frac{\lambda^2 H^2}{2^7 \pi^4} \left\{ \ln^2(a) - \left[\frac{5}{6} + 2\gamma + \ln\left(\frac{H}{2\mu}\right) \right] \ln(a) + \frac{1}{16} - \frac{238}{81} + \frac{13}{54}\pi^2 + \frac{4}{3}\zeta(3) \right\} a^4 \delta^4(x-x'). \quad (69)$$

Note that the nonlocal divergence precisely cancels the nonlocal divergences from the sunset diagram (53) and the snowman diagram (61). As usual, the -- polarity is minus (69).

VII. THE FULLY RENORMALIZED RESULT

To get the full two loop result for the ++ case we add the contributions from the sunset diagram (53), the snowman diagram (61), the counter vertex diagram (64), the field strength renormalization (67) and the two loop mass renormalization (69),

$$M_{2++}^2 = M_{ss++}^2 + M_{sm+}^2 + M_{cv+}^2 + M_{\delta Z+}^2 + M_{\delta m+}^2 \\ = \frac{i\lambda^2}{2^9 \pi^6} \left\{ \frac{aa'}{24} \partial^4 \left[\frac{\ln(\mu^2 \Delta x_{++}^2)}{\Delta x_{++}^2} \right] - H^2 (aa')^2 \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2 \Delta x_{++}^2)}{\Delta x_{++}^2} + \frac{\ln^2(\mu^2 \Delta x_{++}^2)}{4\Delta x_{++}^2} \right] \right. \\ \left. - H^4 (aa')^3 \frac{\ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{++}^2\right)}{\Delta x_{++}^2} + \frac{H^6}{6} (aa')^4 \ln^3\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{++}^2\right) \right\} \\ + \frac{\lambda^2}{2^9 3 \pi^4} a^2 \left\{ -\ln(a) \partial^2 + (2\ln(a) + 1) H a \partial_0 \right\} \delta^4(x-x') \\ + \frac{\lambda^2 H^2}{2^7 \pi^4} \left\{ -\frac{4}{9} \ln^3(a) - \frac{23}{18} \ln^2(a) + \left[\frac{13}{3} + 3\ln\left(\frac{H}{2\mu}\right) - \frac{2}{9}\pi^2 \right] \ln(a) \right\} a^4 \delta^4(x-x') \\ + \frac{\lambda^2 H^2}{2^7 \pi^4} \left\{ \frac{a^{-3}}{81} - \sum_{n=1}^{\infty} \frac{n+5}{(n+1)^3} a^{-(n+1)} + 4 \sum_{n=1}^{\infty} \frac{a^{-(n+2)}}{(n+2)^3} + 4 \sum_{n=1}^{\infty} \frac{a^{-(n+3)}}{n(n+3)^3} \right\} a^4 \delta^4(x-x'). \quad (70)$$

As with the renormalized stress-energy tensor [3], we conjecture that the exponentially falling terms on the last line could be absorbed into a perturbative modification of the initial state. The $--$ polarity is minus the complex conjugate,

$$\begin{aligned}
M_{2--}^2 &= -\left(M_{2++}^2\right)^* \\
&= \frac{i\lambda^2}{2^9\pi^6} \left\{ \frac{aa'}{24} \partial^4 \left[\frac{\ln(\mu^2 \Delta x_{--}^2)}{\Delta x_{--}^2} \right] - H^2(aa')^2 \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2 \Delta x_{--}^2)}{\Delta x_{--}^2} + \frac{\ln^2(\mu^2 \Delta x_{--}^2)}{4\Delta x_{--}^2} \right] \right. \\
&\quad \left. - H^4(aa')^3 \frac{\ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{--}^2\right)}{\Delta x_{--}^2} + \frac{H^6}{6} (aa')^4 \ln^3\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{--}^2\right) \right\} \\
&\quad - \frac{\lambda^2}{2^9 3 \pi^4} a^2 \left\{ -\ln(a) \partial^2 + (2\ln(a) + 1) H a \partial_0 \right\} \delta^4(x-x') \\
&\quad - \frac{\lambda^2 H^2}{2^7 \pi^4} \left\{ -\frac{4}{9} \ln^3(a) - \frac{23}{18} \ln^2(a) + \left[\frac{13}{3} + 3 \ln\left(\frac{H}{2\mu}\right) - \frac{2}{9} \pi^2 \right] \ln(a) \right\} a^4 \delta^4(x-x') \\
&\quad - \frac{\lambda^2 H^2}{2^7 \pi^4} \left\{ \frac{a^{-3}}{81} - \sum_{n=1}^{\infty} \frac{n+5}{(n+1)^3} a^{-(n+1)} + 4 \sum_{n=1}^{\infty} \frac{a^{-(n+2)}}{(n+2)^3} + 4 \sum_{n=1}^{\infty} \frac{a^{-(n+3)}}{n(n+3)^3} \right\} a^4 \delta^4(x-x') . \quad (71)
\end{aligned}$$

As usual, the $+-$ and $-+$ polarities have the opposite sign and lack the local terms,

$$\begin{aligned}
M_{2+-}^2 &= -\frac{i\lambda^2}{2^9\pi^6} \left\{ \frac{aa'}{24} \partial^4 \left[\frac{\ln(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} \right] - H^2(aa')^2 \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} + \frac{\ln^2(\mu^2 \Delta x_{+-}^2)}{4\Delta x_{+-}^2} \right] \right. \\
&\quad \left. - H^4(aa')^3 \frac{\ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{+-}^2\right)}{\Delta x_{+-}^2} + \frac{H^6}{6} (aa')^4 \ln^3\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{+-}^2\right) \right\} \quad (72)
\end{aligned}$$

$$\begin{aligned}
M_{2-+}^2 &= -\frac{i\lambda^2}{2^9\pi^6} \left\{ \frac{aa'}{24} \partial^4 \left[\frac{\ln(\mu^2 \Delta x_{-+}^2)}{\Delta x_{-+}^2} \right] - H^2(aa')^2 \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2 \Delta x_{-+}^2)}{\Delta x_{-+}^2} + \frac{\ln^2(\mu^2 \Delta x_{-+}^2)}{4\Delta x_{-+}^2} \right] \right. \\
&\quad \left. - H^4(aa')^3 \frac{\ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{-+}^2\right)}{\Delta x_{-+}^2} + \frac{H^6}{6} (aa')^4 \ln^3\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{-+}^2\right) \right\} \quad (73)
\end{aligned}$$

VIII. DISCUSSION

We have computed and fully renormalized the four Schwinger-Keldysh self-mass-squared's at one and two loop orders for massless, minimally coupled φ^4 (2) on a locally de Sitter background (1). The next step is using our results to study the quantum-corrected, effective field equations at linearized order (7). The term “ $M^2(x; x')$ ” called for in (7) is the sum of $M_{++}^2(x; x')$ and $M_{+-}^2(x; x')$. The analysis — which will be presented in a subsequent paper — is very similar to that already done for de Sitter photons using the one loop vacuum polarization [87] and for de Sitter fermions using the one loop self-energy [88].

There are two interesting issues to investigate. The first is stability. Secular effects in this system are driven by inflationary particle production which forces the scalar up its φ^4 potential. One would expect this to generate a *positive* mass-squared, and it is significant that our one loop result (41) is positive. The two loop result (70) is a little too complicated to interpret without a detailed calculation of the type proposed, but we expect no problems with stability.

The second interesting issue is an apparent paradox in the stochastic solution of this model which was given in 1994 by Starobinskiĭ and Yokoyama [89]. As we noted in the introduction, this model is stable for two reasons. First, as inflationary particle production forces the scalar up its potential the classical force pushes it back down. Second, the curvature of the potential ought to induce a positive mass-squared which should reduce the inflationary particle production. However, the solution of Starobinskiĭ and Yokoyama implements only the first effect. Although it would be possible to implement the second effect as well within the context of Starobinskiĭ's stochastic technique [90], including just the first effect gives perfect agreement at leading logarithm order as high as this has been checked. The question is why should this be so? A possible answer is that the nonzero self-mass-squared remains always down by one factor of $\ln(a)$. Whereas the leading logarithms of the energy density give a series in $\lambda \ln^2(a)$,

$$\rho|_{\text{leading}} = H^4 \sum_{\ell=2}^{\infty} \rho_{\ell} \left(\lambda \ln^2(a) \right)^{\ell-1}, \quad (74)$$

it may be that the self-mass-squared behaves as such a series times $\lambda \ln(a)$,

$$M^2(x; x')|_{\text{leading}} = H^2 a^4 \delta^4(x - x') \times \lambda \ln(a) \sum_{\ell=1}^{\infty} m_{\ell} \left(\lambda \ln^2(a) \right)^{\ell-1}. \quad (75)$$

It is worth noting that, had gravity been quantized, there would have been similar corrections from graviton exchange. While the physics of the pure scalar model pretty much guarantees that a positive self-mass-squared must be induced, there is no such argument for gravity. If the induced self-mass-squared turns out to be tachyonic it might have important implications for scalar-driven inflation. Further, there is no reason the gravitational process should ever stop. The graviton should not experience a classical force since it has no non-derivative potential. Nor should the graviton ever develop a mass. This might be significant for scalar-driven models which inflate for a very large number of e-foldings.

We close with a comment on accuracy. Two loop computations are generally intricate enough that this becomes an issue. In this regard it is worth noting that three distinct

diagrams conspire to cancel the overlapping divergences. This is significant because there are no overlapping divergences at this order in massless φ^4 on flat space.

It is also a significant correspondence check that we get the same results for $\delta\lambda$ (at one loop) and δZ (at two loops) as for massless φ^4 in flat space. This had to be the case because the divergences these counterterms absorb are formally logarithmic. Because geometric corrections make divergences more finite, any geometric corrections to $\delta\lambda$ and δZ must be finite. The same is not the case for δm^2 because the divergences it cancels are formally quadratic. Indeed, δm^2 vanishes for massless φ^4 in flat space whereas we found logarithmically divergent corrections proportional to H^2 at both one and two loop orders.

IX. APPENDIX

A. Partial Integration Identities

In almost all cases derivatives of the conformal coordinate intervals $\Delta x_{\pm\pm}^2(x; x')$ can be taken generically, without regard to polarity and without worrying about delta functions,

$$\partial^2 \left(\frac{1}{\Delta x_{\pm\pm}^{n-m\epsilon}} \right) = \frac{(n-m\epsilon)(n-2-(m-1)\epsilon)}{\Delta x_{\pm\pm}^{n+2-m\epsilon}} \quad (n \neq 2 \text{ or } m \neq 1). \quad (76)$$

The exception is $1/\Delta x^{2-\epsilon}$,

$$\partial^2 \left(\frac{1}{\Delta x_{++}^{2-\epsilon}} \right) = \frac{i4\pi^{2-\frac{\epsilon}{2}}}{\Gamma(1-\frac{\epsilon}{2})} \delta^D(x-x') = -\partial^2 \left(\frac{1}{\Delta x_{--}^{2-\epsilon}} \right) \quad , \quad \partial^2 \left(\frac{1}{\Delta x_{+-}^{2-\epsilon}} \right) = 0 = \partial^2 \left(\frac{1}{\Delta x_{-+}^{2-\epsilon}} \right). \quad (77)$$

These identities allow one to extract more and more derivatives of inverse powers of $\Delta x_{\pm\pm}^2$ until the result is integrable in $D = 4$ dimensions. For $m \neq 1$ we have,

$$\frac{1}{\Delta x_{\pm\pm}^{4-m\epsilon}} = \frac{\partial^2}{(2-m\epsilon)(-(m-1)\epsilon)} \left(\frac{1}{\Delta x_{\pm\pm}^{2-m\epsilon}} \right), \quad (78)$$

$$\frac{1}{\Delta x_{\pm\pm}^{6-m\epsilon}} = \frac{\partial^4}{(4-m\epsilon)(2-m\epsilon)(2-(m-1)\epsilon)(-(m-1)\epsilon)} \left(\frac{1}{\Delta x_{\pm\pm}^{2-m\epsilon}} \right). \quad (79)$$

At this stage one adds zero in the form of the appropriate identity (77) and then expands the nonlocal term in powers of ϵ . For example, the $++$ term gives,

$$\partial^2 \left(\frac{1}{\Delta x_{++}^{2-m\epsilon}} \right) = \partial^2 \left(\frac{1}{\Delta x_{++}^{2-m\epsilon}} - \frac{\mu^{-(m-1)\epsilon}}{\Delta x_{++}^{2-\epsilon}} \right) + \frac{i4\pi^{2-\frac{\epsilon}{2}}\mu^{-(m-1)\epsilon}}{\Gamma(1-\frac{\epsilon}{2})} \delta^D(x-x'), \quad (80)$$

$$\begin{aligned}
&= + \frac{i4\pi^{2-\frac{\epsilon}{2}}\mu^{-(m-1)\epsilon}}{\Gamma(1-\frac{\epsilon}{2})} \delta^D(x-x') \\
&\quad + \mu^{-m\epsilon} \partial^2 \left(\frac{\frac{(m-1)}{2}\epsilon \ln(\mu^2 \Delta x_{++}^2) + \frac{(m^2-1)}{8}\epsilon^2 \ln^2(\mu^2 \Delta x_{++}^2) + O(\epsilon^3)}{\Delta x_{++}^2} \right). \quad (81)
\end{aligned}$$

The cases we require are,

$$\begin{aligned}
\frac{1}{\Delta x_{++}^{4-2\epsilon}} &= \frac{\mu^{-\epsilon}}{(2-2\epsilon)(-\epsilon)} \frac{i4\pi^{2-\frac{\epsilon}{2}}}{\Gamma(1-\frac{\epsilon}{2})} \delta^D(x-x') \\
&\quad + \frac{\mu^{-2\epsilon} \partial^2}{(2-2\epsilon)(-\epsilon)} \left(\frac{\frac{1}{2}\epsilon \ln(\mu^2 \Delta x_{++}^2) + \frac{3}{8}\epsilon^2 \ln^2(\mu^2 \Delta x_{++}^2) + O(\epsilon^3)}{\Delta x_{++}^2} \right), \quad (82)
\end{aligned}$$

$$\begin{aligned}
\frac{1}{\Delta x_{++}^{4-3\epsilon}} &= \frac{\mu^{-2\epsilon}}{(2-3\epsilon)(-2\epsilon)} \frac{i4\pi^{2-\frac{\epsilon}{2}}}{\Gamma(1-\frac{\epsilon}{2})} \delta^D(x-x') \\
&\quad + \frac{\mu^{-3\epsilon} \partial^2}{(2-3\epsilon)(-2\epsilon)} \left(\frac{\epsilon \ln(\mu^2 \Delta x_{++}^2) + \epsilon^2 \ln^2(\mu^2 \Delta x_{++}^2) + O(\epsilon^3)}{\Delta x_{++}^2} \right), \quad (83)
\end{aligned}$$

$$\begin{aligned}
\frac{1}{\Delta x_{++}^{6-3\epsilon}} &= \frac{\mu^{-2\epsilon}}{(4-3\epsilon)(2-3\epsilon)(2-2\epsilon)(-2\epsilon)} \frac{i4\pi^{2-\frac{\epsilon}{2}}}{\Gamma(1-\frac{\epsilon}{2})} \partial^2 \delta^D(x-x') \\
&\quad + \frac{\mu^{-3\epsilon} \partial^4}{(4-3\epsilon)(2-3\epsilon)(2-2\epsilon)(-2\epsilon)} \left(\frac{\epsilon \ln(\mu^2 \Delta x_{++}^2) + O(\epsilon^2)}{\Delta x_{++}^2} \right). \quad (84)
\end{aligned}$$

The -- variations differ only in the sign of the delta function term,

$$\begin{aligned}
\frac{1}{\Delta x_{--}^{4-2\epsilon}} &= - \frac{\mu^{-\epsilon}}{(2-2\epsilon)(-\epsilon)} \frac{i4\pi^{2-\frac{\epsilon}{2}}}{\Gamma(1-\frac{\epsilon}{2})} \delta^D(x-x') \\
&\quad + \frac{\mu^{-2\epsilon} \partial^2}{(2-2\epsilon)(-\epsilon)} \left(\frac{\frac{1}{2}\epsilon \ln(\mu^2 \Delta x_{--}^2) + \frac{3}{8}\epsilon^2 \ln^2(\mu^2 \Delta x_{--}^2) + O(\epsilon^3)}{\Delta x_{--}^2} \right), \quad (85)
\end{aligned}$$

$$\begin{aligned}
\frac{1}{\Delta x_{--}^{4-3\epsilon}} &= - \frac{\mu^{-2\epsilon}}{(2-3\epsilon)(-2\epsilon)} \frac{i4\pi^{2-\frac{\epsilon}{2}}}{\Gamma(1-\frac{\epsilon}{2})} \delta^D(x-x') \\
&\quad + \frac{\mu^{-3\epsilon} \partial^2}{(2-3\epsilon)(-2\epsilon)} \left(\frac{\epsilon \ln(\mu^2 \Delta x_{--}^2) + \epsilon^2 \ln^2(\mu^2 \Delta x_{--}^2) + O(\epsilon^3)}{\Delta x_{--}^2} \right), \quad (86)
\end{aligned}$$

$$\begin{aligned}
\frac{1}{\Delta x_{--}^{6-3\epsilon}} &= - \frac{\mu^{-2\epsilon}}{(4-3\epsilon)(2-3\epsilon)(2-2\epsilon)(-2\epsilon)} \frac{i4\pi^{2-\frac{\epsilon}{2}}}{\Gamma(1-\frac{\epsilon}{2})} \partial^2 \delta^D(x-x') \\
&\quad + \frac{\mu^{-3\epsilon} \partial^4}{(4-3\epsilon)(2-3\epsilon)(2-2\epsilon)(-2\epsilon)} \left(\frac{\epsilon \ln(\mu^2 \Delta x_{--}^2) + O(\epsilon^2)}{\Delta x_{--}^2} \right). \quad (87)
\end{aligned}$$

And the mixed terms have no delta functions,

$$\frac{1}{\Delta x_{+-}^{4-2\epsilon}} = \frac{\mu^{-2\epsilon} \partial^2}{(2-2\epsilon)(-\epsilon)} \left(\frac{\frac{1}{2}\epsilon \ln(\mu^2 \Delta x_{+-}^2) + \frac{3}{8}\epsilon^2 \ln^2(\mu^2 \Delta x_{+-}^2) + O(\epsilon^3)}{\Delta x_{+-}^2} \right), \quad (88)$$

$$\frac{1}{\Delta x_{+-}^{4-3\epsilon}} = \frac{\mu^{-3\epsilon} \partial^2}{(2-3\epsilon)(-2\epsilon)} \left(\frac{\epsilon \ln(\mu^2 \Delta x_{+-}^2) + \epsilon^2 \ln^2(\mu^2 \Delta x_{+-}^2) + O(\epsilon^3)}{\Delta x_{+-}^2} \right), \quad (89)$$

$$\frac{1}{\Delta x_{+-}^{6-3\epsilon}} = \frac{\mu^{-3\epsilon} \partial^4}{(4-3\epsilon)(2-3\epsilon)(2-2\epsilon)(-2\epsilon)} \left(\frac{\epsilon \ln(\mu^2 \Delta x_{+-}^2) + O(\epsilon^2)}{\Delta x_{+-}^2} \right). \quad (90)$$

Replacing $+-$ with $-+$ gives the final case.

B. The Square of a Propagator

When the square of a propagator multiplies regular expressions we first decompose it according to (34) and then take ϵ to zero in all but the A^2 term,

$$(i\Delta(x; x'))^2 = (A(x; x'))^2 + 2A(x; x')(B(x; x') + C(x; x')) + (B(x; x') + C(x; x'))^2, \quad (91)$$

$$\longrightarrow \frac{\Gamma^2(1-\frac{\epsilon}{2})(aa')^{-2+\epsilon}}{2^4\pi^{4-\epsilon}} \frac{1}{\Delta x^{4-2\epsilon}} - \frac{H^2}{2^4\pi^4} \frac{\ln(\frac{\sqrt{e}}{4} H^2 \Delta x^2)}{aa' \Delta x^2} + \frac{H^4}{2^6\pi^4} \ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x^2\right). \quad (92)$$

We then employ the appropriate identity — (82), (85) or (88) — and take $\epsilon = 0$ in the nonlocal part. The squares of the $+-$ and $-+$ propagators are finite,

$$(i\Delta_{+-})^2 \longrightarrow -\frac{(aa')^{-2}}{2^6\pi^4} \partial^2 \left(\frac{\ln(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} \right) - \frac{H^2}{2^4\pi^4} \frac{\ln(\frac{\sqrt{e}}{4} H^2 \Delta x_{+-}^2)}{aa' \Delta x_{+-}^2} + \frac{H^4}{2^6\pi^4} \ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{+-}^2\right), \quad (93)$$

$$(i\Delta_{-+})^2 \longrightarrow -\frac{(aa')^{-2}}{2^6\pi^4} \partial^2 \left(\frac{\ln(\mu^2 \Delta x_{-+}^2)}{\Delta x_{-+}^2} \right) - \frac{H^2}{2^4\pi^4} \frac{\ln(\frac{\sqrt{e}}{4} H^2 \Delta x_{-+}^2)}{aa' \Delta x_{-+}^2} + \frac{H^4}{2^6\pi^4} \ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{-+}^2\right). \quad (94)$$

The square of the $++$ propagator includes a local divergent term in addition to the corresponding $++$ finite terms,

$$(i\Delta_{++}(x; x'))^2 \longrightarrow -\frac{i\mu^{-\epsilon}}{2^3\pi^{2-\frac{\epsilon}{2}}} \frac{\Gamma(1-\frac{\epsilon}{2})}{(1-\epsilon)\epsilon} a^{-4+2\epsilon} \delta^D(x-x') - \frac{(aa')^{-2}}{2^6\pi^4} \partial^2 \left(\frac{\ln(\mu^2 \Delta x_{++}^2)}{\Delta x_{++}^2} \right) - \frac{H^2}{2^4\pi^4} \frac{\ln(\frac{\sqrt{e}}{4} H^2 \Delta x_{++}^2)}{aa' \Delta x_{++}^2} + \frac{H^4}{2^6\pi^4} \ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{++}^2\right). \quad (95)$$

The square of the $--$ propagator is the complex conjugate,

$$(i\Delta_{--}(x; x'))^2 \longrightarrow \frac{i\mu^{-\epsilon}}{2^3\pi^{2-\frac{\epsilon}{2}}} \frac{\Gamma(1-\frac{\epsilon}{2})}{(1-\epsilon)\epsilon} a^{-4+2\epsilon} \delta^D(x-x') - \frac{(aa')^{-2}}{2^6\pi^4} \partial^2 \left(\frac{\ln(\mu^2 \Delta x_{--}^2)}{\Delta x_{--}^2} \right) - \frac{H^2}{2^4\pi^4} \frac{\ln(\frac{\sqrt{e}}{4} H^2 \Delta x_{--}^2)}{aa' \Delta x_{--}^2} + \frac{H^4}{2^6\pi^4} \ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{--}^2\right). \quad (96)$$

C. The Cube of a Propagator

When the cube of a propagator multiplies a regular expression we again decompose it according to (34) but now we must retain $\epsilon \neq 0$ in the A^3 and $3A^2B$ terms,

$$(i\Delta(x; x'))^3 = A^3 + 3A^2(B+C) + 3A(B+C)^2 + (B+C)^3, \quad (97)$$

$$\begin{aligned} \longrightarrow & \frac{\Gamma^3(1-\frac{\epsilon}{2}) (aa')^{-3+\frac{3}{2}\epsilon}}{2^6\pi^{6-\frac{3}{2}\epsilon} \Delta x^{6-3\epsilon}} + \frac{3H^{2-\epsilon}\Gamma(1-\frac{\epsilon}{2})\Gamma(2-\epsilon)}{2^{7-\epsilon}\pi^{6-\frac{3}{2}\epsilon}(aa')^{2-\epsilon}} \left\{ -\frac{2\Gamma(3-\frac{\epsilon}{2})\Gamma(2-\frac{\epsilon}{2})}{\epsilon} \frac{(\frac{H^2aa'}{4})^{\frac{\epsilon}{2}}}{\Delta x^{4-3\epsilon}} \right. \\ & \left. + \left[\frac{\pi \cot(\frac{\pi\epsilon}{2}) + \ln(aa')}{\Delta x^{4-2\epsilon}} \right] \right\} + \frac{3H^4 \ln^2(\frac{\sqrt{e}}{4}H^2\Delta x^2)}{2^8\pi^6 aa' \Delta x^2} - \frac{H^6}{2^9\pi^6} \ln^3(\frac{\sqrt{e}}{4}H^2\Delta x^2). \quad (98) \end{aligned}$$

The A^3 term is easy to read off from the identities (84), (87) and (90). For example, the $+-$ case gives,

$$\frac{\Gamma^3(1-\frac{\epsilon}{2}) (aa')^{-3+\frac{3}{2}\epsilon}}{2^6\pi^{6-\frac{3}{2}\epsilon} \Delta x_{+-}^{6-3\epsilon}} \longrightarrow -\frac{(aa')^{-3}}{2^{11}\pi^6} \partial^4 \left[\frac{\ln(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} \right]. \quad (99)$$

The $3A^2B$ term is more complicated on account of the $1/\epsilon$ divergence in its prefactors. One must first partially integrate and multiply before taking $\epsilon = 0$ in the nonlocal term. For the $+-$ case we have,

$$\begin{aligned} 3(A^2B)_{+-} = & \frac{3H^{2-\epsilon}(aa')^{-2+\epsilon}\mu^{-2\epsilon}}{2^{7-\epsilon}\pi^{6-\frac{3}{2}\epsilon}} \Gamma(1-\frac{\epsilon}{2}) \\ & \times \left\{ \frac{(1-\frac{\epsilon}{2})(1-\frac{\epsilon}{4})\Gamma^2(1-\frac{\epsilon}{2})}{2(1-\frac{3}{2}\epsilon)\epsilon} \left(\frac{H^2aa'}{4\mu^2} \right)^{\frac{\epsilon}{2}} \partial^2 \left[\frac{\ln(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} + \epsilon \frac{\ln^2(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} \right] \right. \\ & \left. - \frac{\Gamma(1-\epsilon)}{4} \left[\pi \cot\left(\frac{\pi\epsilon}{2}\right) + \ln(aa') \right] \partial^2 \left[\frac{\ln(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} + \frac{3}{4}\epsilon \frac{\ln^2(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} \right] \right\}. \quad (100) \end{aligned}$$

At this point we digress to recall some basic series expansions,

$$\Gamma(1-\epsilon) = 1 + \gamma\epsilon + \left[\frac{\gamma^2}{2} + \frac{\pi^2}{12} \right] \epsilon^2 + O(\epsilon^3) \quad , \quad \pi \cot\left(\frac{\pi\epsilon}{2}\right) = \frac{2}{\epsilon} \left[1 - \frac{\pi^2}{12} \epsilon^2 + O(\epsilon^4) \right]. \quad (101)$$

It follows that the various prefactors in (100) are,

$$\frac{(1-\frac{\epsilon}{2})(1-\frac{\epsilon}{4})\Gamma^2(1-\frac{\epsilon}{2})}{2(1-\frac{3}{2}\epsilon)\epsilon} = \frac{1}{2\epsilon} + \frac{\gamma}{2} + \frac{3}{8} + O(\epsilon), \quad (102)$$

$$-\frac{\Gamma(1-\epsilon)}{4} \pi \cot\left(\frac{\pi\epsilon}{2}\right) = -\frac{1}{2\epsilon} - \frac{\gamma}{2} + O(\epsilon), \quad (103)$$

$$-\frac{\Gamma(1-\epsilon)}{4} = -\frac{1}{4} + O(\epsilon). \quad (104)$$

Making use of these expansions in (100) gives,

$$3(A^2B)_{+-} \longrightarrow \frac{3H^2(aa')^{-2}}{2^8\pi^6} \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} + \frac{1}{4} \frac{\ln^2(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} \right]. \quad (105)$$

Employing (99) and (105) in (98) gives

$$(i\Delta_{+-}(x; x'))^3 \rightarrow -\frac{(aa')^{-3}}{2^{11}\pi^6} \partial^4 \left[\frac{\ln(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} \right] + \frac{3H^2(aa')^{-2}}{2^8\pi^6} \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} \right. \\ \left. + \frac{1}{4} \frac{\ln^2(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} \right] + \frac{3H^4 \ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{+-}^2\right)}{2^8\pi^6 aa' \Delta x_{+-}^2} - \frac{H^6}{2^9\pi^6} \ln^3\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{+-}^2\right). \quad (106)$$

The $-+$ case follows by complex conjugation,

$$(i\Delta_{-+}(x; x'))^3 \rightarrow -\frac{(aa')^{-3}}{2^{11}\pi^6} \partial^4 \left[\frac{\ln(\mu^2 \Delta x_{-+}^2)}{\Delta x_{-+}^2} \right] + \frac{3H^2(aa')^{-2}}{2^8\pi^6} \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2 \Delta x_{-+}^2)}{\Delta x_{-+}^2} \right. \\ \left. + \frac{1}{4} \frac{\ln^2(\mu^2 \Delta x_{-+}^2)}{\Delta x_{-+}^2} \right] + \frac{3H^4 \ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{-+}^2\right)}{2^8\pi^6 aa' \Delta x_{-+}^2} - \frac{H^6}{2^9\pi^6} \ln^3\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{-+}^2\right). \quad (107)$$

The cube of the $++$ propagator includes a local divergent term in addition to the corresponding $++$ finite terms,

$$(i\Delta_{++}(x; x'))^3 \rightarrow -\frac{i\mu^{-2\epsilon}\Gamma^2(1-\frac{\epsilon}{2})}{2^9\pi^{4-\epsilon}} \frac{(aa')^{-3+\frac{3}{2}\epsilon}\partial^2\delta^D(x-x')}{(1-\frac{3}{2}\epsilon)(1-\epsilon)(1-\frac{3}{4}\epsilon)\epsilon} + \frac{i3H^{2-\epsilon}\mu^{-\epsilon}a^{-4+2\epsilon}}{2^{6-\epsilon}\pi^{4-\epsilon}} \left\{ \right. \\ \left. \frac{(1-\frac{\epsilon}{2})(1-\frac{\epsilon}{4})\Gamma^2(1-\frac{\epsilon}{2})}{(1-\frac{3}{2}\epsilon)\epsilon^2} \left(\frac{Ha}{2\mu}\right)^\epsilon - \frac{\Gamma(1-\epsilon)}{\epsilon} \left[\pi \cot\left(\frac{\pi\epsilon}{2}\right) + 2\ln(a) \right] \right\} \delta^D(x-x') \\ - \frac{(aa')^{-3}}{2^{11}\pi^6} \partial^4 \left[\frac{\ln(\mu^2 \Delta x_{++}^2)}{\Delta x_{++}^2} \right] + \frac{3H^2(aa')^{-2}}{2^8\pi^6} \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2 \Delta x_{++}^2)}{\Delta x_{++}^2} + \frac{1}{4} \frac{\ln^2(\mu^2 \Delta x_{++}^2)}{\Delta x_{++}^2} \right] \\ + \frac{3H^4 \ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{++}^2\right)}{2^8\pi^6 aa' \Delta x_{++}^2} - \frac{H^6}{2^9\pi^6} \ln^3\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{++}^2\right). \quad (108)$$

The cube of the $--$ propagator is the complex conjugate,

$$(i\Delta_{--}(x; x'))^3 \rightarrow \frac{i\mu^{-2\epsilon}\Gamma^2(1-\frac{\epsilon}{2})}{2^9\pi^{4-\epsilon}} \frac{(aa')^{-3+\frac{3}{2}\epsilon}\partial^2\delta^D(x-x')}{(1-\frac{3}{2}\epsilon)(1-\epsilon)(1-\frac{3}{4}\epsilon)\epsilon} - \frac{i3H^{2-\epsilon}\mu^{-\epsilon}a^{-4+2\epsilon}}{2^{6-\epsilon}\pi^{4-\epsilon}} \left\{ \right. \\ \left. \frac{(1-\frac{\epsilon}{2})(1-\frac{\epsilon}{4})\Gamma^2(1-\frac{\epsilon}{2})}{(1-\frac{3}{2}\epsilon)\epsilon^2} \left(\frac{Ha}{2\mu}\right)^\epsilon - \frac{\Gamma(1-\epsilon)}{\epsilon} \left[\pi \cot\left(\frac{\pi\epsilon}{2}\right) + 2\ln(a) \right] \right\} \delta^D(x-x') \\ - \frac{(aa')^{-3}}{2^{11}\pi^6} \partial^4 \left[\frac{\ln(\mu^2 \Delta x_{--}^2)}{\Delta x_{--}^2} \right] + \frac{3H^2(aa')^{-2}}{2^8\pi^6} \partial^2 \left[\ln\left(\frac{He^{\frac{3}{4}}}{2\mu}\right) \frac{\ln(\mu^2 \Delta x_{--}^2)}{\Delta x_{--}^2} + \frac{1}{4} \frac{\ln^2(\mu^2 \Delta x_{--}^2)}{\Delta x_{--}^2} \right] \\ + \frac{3H^4 \ln^2\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{--}^2\right)}{2^8\pi^6 aa' \Delta x_{--}^2} - \frac{H^6}{2^9\pi^6} \ln^3\left(\frac{\sqrt{e}}{4} H^2 \Delta x_{--}^2\right). \quad (109)$$

D. Evaluating the Integrals for the Snowman Diagram

The first integral we need in equation (59) is,

$$I_1 \equiv -\partial_0^2 \int d^4x'' a''^2 \ln(a'') \left[\frac{\ln(\mu^2 \Delta x_{++}^2)}{\Delta x_{++}^2} - \frac{\ln(\mu^2 \Delta x_{+-}^2)}{\Delta x_{+-}^2} \right]. \quad (110)$$

The first step is to make use of the identity,

$$\frac{\ln(\mu^2 \Delta x^2)}{\Delta x^2} = \frac{\partial^2}{8} \left[\ln^2(\mu^2 \Delta x^2) - 2 \ln(\mu^2 \Delta x^2) \right]. \quad (111)$$

Because the derivative is with respect to x^μ we can move it outside the integration over $x^{\mu\mu}$.

The remaining integrand possesses only logarithmic singularities.

We define the coordinate separations,

$$\Delta\eta \equiv \eta - \eta'' \quad , \quad r \equiv \|\vec{x} - \vec{x}''\|. \quad (112)$$

The $++$ and $+-$ terms cancel for $\Delta\eta < 0$ so we can restrict the integration to $\Delta\eta > 0$. In this case the logarithms can be expanded as,

$$\ln[\mu^2 \Delta x_{\pm}^2] = \ln[\mu^2(\Delta\eta^2 - r^2)] \pm i\pi\theta(\Delta\eta^2 - r^2), \quad (113)$$

We then perform the angular and radial integrations and act the first three derivatives,

$$\begin{aligned} I_1 &= i2\pi^2 \partial_0^4 \int_{\eta_i}^{\eta} d\eta'' a''^2 \ln(a'') \int_0^{\Delta\eta} dr r^2 \left(\ln[\mu^2(\Delta\eta^2 - r^2)] - 1 \right) \\ &= i2\pi^2 \partial_0^4 \int_{\eta_i}^{\eta} d\eta'' a''^2 \ln(a'') \Delta\eta^3 \left[\frac{2}{3} \ln(2\mu\Delta\eta) - \frac{11}{9} \right] \\ &= i8\pi^2 \partial_0 \int_{\eta_i}^{\eta} d\eta'' a''^2 \ln(a'') \ln(2\mu\Delta\eta). \end{aligned} \quad (114)$$

The lower limit $\eta_i = -H^{-1}$ ($t'' = 0$) derives from the fact that the state is released in free Bunch-Davies vacuum at this instant.

To perform the remaining temporal integration first change variables from η'' to $a'' = -1/(H\eta'')$ and make the corresponding change in the derivative,

$$I_1 = i8\pi^2 a^2 \frac{\partial}{\partial a} \int_1^a da'' \ln(a'') \ln \left[\frac{2\mu}{H} \left(\frac{1}{a''} - \frac{1}{a} \right) \right]. \quad (115)$$

Then break up the second logarithm and act the derivative on the two nonsingular terms,

$$I_1 = -i8\pi^2 a^2 \left[\ln^2(a) - \ln\left(\frac{2\mu}{H}\right) \ln(a) - \frac{\partial}{\partial a} \int_1^a da'' \ln(a'') \ln \left(1 - \frac{a''}{a} \right) \right]. \quad (116)$$

Now substitute the expansion,

$$\ln \left(1 - \frac{a''}{a} \right) = - \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{a''}{a} \right)^n, \quad (117)$$

and integrate termwise,

$$\int_1^a da'' \ln(a'') \ln \left(1 - \frac{a''}{a} \right) = -a \left\{ \ln(a) - 2 + \frac{\pi^2}{6} + \sum_{n=1}^{\infty} \frac{a^{-(n+1)}}{n(n+1)^2} \right\}. \quad (118)$$

Combining terms gives the following final result,

$$I_1 = -i8\pi^2 a^2 \left[\ln^2(a) - \ln\left(\frac{2\mu}{H}\right) \ln(a) + \ln(a) - 1 + \frac{\pi^2}{6} - \sum_{n=1}^{\infty} \frac{a^{-(n+1)}}{(n+1)^2} \right]. \quad (119)$$

The second integral in (59) is,

$$I_2 \equiv \int d^4 x'' a''^3 \ln(a'') \left[\frac{\ln\left(\frac{H^2 \sqrt{e}}{4} \Delta x_{++}^2\right)}{\Delta x_{++}^2} - \frac{\ln\left(\frac{H^2 \sqrt{e}}{4} \Delta x_{+-}^2\right)}{\Delta x_{+-}^2} \right]. \quad (120)$$

Proceeding as before we extract a derivative with identity (111) and then exploit (113) to obtain,

$$\begin{aligned} I_2 &= i\frac{\pi}{2} \partial^2 \int d^4 x'' a''^3 \ln(a'') \theta(\Delta\eta^2 - r^2) \ln\left(\frac{H^2}{4\sqrt{e}}(\Delta\eta^2 - r^2)\right) \\ &= i2\pi^2 (-\partial_0^2) \int_{\eta_i}^{\eta} d\eta'' a''^3 \ln(a'') \int_0^{\Delta\eta} dr r^2 \ln\left(\frac{H^2}{4\sqrt{e}}(\Delta\eta^2 - r^2)\right). \end{aligned} \quad (121)$$

Now perform the angular integrations, make the change of variable $r \equiv \Delta\eta z$, act the temporal derivatives and change variables from η'' to a'' ,

$$\begin{aligned} I_2 &= -i2\pi^2 \int_{\eta_i}^{\eta} d\eta'' a''^3 \ln(a'') \int_0^1 dz z^2 \partial_0^2 \left[\Delta\eta^3 \ln\left(\frac{H^2}{4\sqrt{e}} \Delta\eta^2 (1 - z^2)\right) \right] \\ &= -i2\pi^2 \int_{\eta_i}^{\eta} d\eta'' a''^3 \ln(a'') \Delta\eta \int_0^1 dz z^2 \left\{ 6 \ln\left[\frac{H^2}{4} \Delta\eta^2 (1 - z^2)\right] + 7 \right\} \\ &= -\frac{i2\pi^2}{H^2} \int_1^a da'' \ln(a'') \left(1 - \frac{a''}{a}\right) \left\{ 4 \ln\left(\frac{1}{a''} - \frac{1}{a}\right) - 3 \right\}. \end{aligned} \quad (122)$$

A small rearrangement gives,

$$I_2 = \frac{i2\pi^2}{H^2} \left\{ \int_1^a da'' \left(1 - \frac{a''}{a}\right) \left[3 \ln(a'') + 4 \ln^2(a'') \right] - 4 \int_1^a da'' \left(1 - \frac{a''}{a}\right) \ln(a'') \ln\left(1 - \frac{a''}{a}\right) \right\}. \quad (123)$$

The only non-trivial integral left in (123) can be evaluated using (117)

$$\int_1^a da'' \left(1 - \frac{a''}{a}\right) \ln(a'') \ln\left(1 - \frac{a''}{a}\right) = -a \left\{ \frac{1}{4} \ln(a) - 1 + \frac{\pi^2}{12} + \sum_{n=1}^{\infty} \left[\frac{a^{-(n+1)}}{n(n+1)^2} - \frac{a^{-(n+2)}}{n(n+2)^2} \right] \right\}. \quad (124)$$

The final result for I_2 is therefore,

$$I_2 = \frac{i2\pi^2 a}{H^2} \left\{ 2 \ln^2(a) - \frac{7}{2} \ln(a) + \frac{3}{4} + \frac{\pi^2}{3} - 5a^{-1} + \frac{1}{4} a^{-2} + 4 \sum_{n=1}^{\infty} \left[\frac{a^{-(n+1)}}{n(n+1)^2} - \frac{a^{-(n+2)}}{n(n+2)^2} \right] \right\}. \quad (125)$$

The initial reduction of the last integral in (59) is familiar from I_1 and I_2 ,

$$I_3 \equiv \int d^4 x'' a''^4 \ln(a'') \left[\ln^2\left(\frac{H^2 \sqrt{e}}{4} \Delta x_{++}^2\right) - \ln^2\left(\frac{H^2 \sqrt{e}}{4} \Delta x_{+-}^2\right) \right]$$

$$\begin{aligned}
&= i4\pi \int d^4x'' a''^4 \ln(a'') \theta(\Delta\eta^2 - r^2) \ln\left(\frac{H^2\sqrt{e}}{4}(\Delta\eta^2 - r^2)\right) \\
&= i16\pi^2 \int_0^1 dz z^2 \int_{\eta_i}^{\eta} d\eta'' a''^4 \ln(a'') \Delta\eta^3 \ln\left(\frac{H^2\sqrt{e}}{4}\Delta\eta^2(1-z^2)\right) \\
&= \frac{i16\pi^2}{H^4} \int_0^1 dz z^2 \int_1^a \frac{da''}{a''} \ln(a'') \left(1 - \frac{a''}{a}\right)^3 \ln\left(\frac{\sqrt{e}}{4}\left(\frac{1}{a''} - \frac{1}{a}\right)^2 (1-z^2)\right) \\
&= \frac{i16\pi^2}{H^4} \int_1^a \frac{da''}{a''} \ln(a'') \left(1 - \frac{a''}{a}\right)^3 \left\{ \frac{2}{3} \ln\left(\frac{1}{a''} - \frac{1}{a}\right) - \frac{13}{18} \right\}. \tag{126}
\end{aligned}$$

At this stage it is best to integrate by parts on the term $\ln(a'')/a'' = \partial/\partial a''(\ln^2(a'')/2)$,

$$\begin{aligned}
I_3 = \frac{i16\pi^2}{H^4} \left\{ - \int_1^a da'' \left(1 - \frac{a''}{a}\right)^2 \ln^2(a'') \left[\frac{1}{a} \left(\frac{13}{12} + \ln(a'') \right) - \frac{1}{3a''} \right] \right. \\
\left. + \frac{1}{a} \int_1^a da'' \left(1 - \frac{a''}{a}\right)^2 \ln^2(a'') \ln\left(1 - \frac{a''}{a}\right) \right\}. \tag{127}
\end{aligned}$$

The first term is straightforward and the final term can be evaluated using (117),

$$\begin{aligned}
\int_1^a da'' \left(1 - \frac{a''}{a}\right)^2 \ln^2(a'') \ln\left(1 - \frac{a''}{a}\right) = a \left\{ -\frac{80}{27} + \frac{11}{54}\pi^2 + \frac{2}{3}\zeta(3) + \left(\frac{71}{54} - \frac{\pi^2}{9}\right) \ln(a) \right. \\
\left. - \frac{1}{9} \ln^2(a) + 2 \sum_{n=1}^{\infty} \left[\frac{a^{-(n+1)}}{n(n+1)^3} - \frac{2a^{-(n+2)}}{n(n+2)^3} + \frac{a^{-(n+3)}}{n(n+3)^3} \right] \right\}. \tag{128}
\end{aligned}$$

Substituting (128) in (127) and adding everything up gives,

$$\begin{aligned}
I_3 = i \frac{8\pi^2}{H^4} \left\{ -\frac{385}{324} + \frac{11}{27}\pi^2 + \frac{4}{3}\zeta(3) + \frac{a^{-3}}{81} + \frac{a^{-2}}{4} - 5a^{-1} - \left(\frac{11}{6} + \frac{2\pi^2}{9}\right) \ln(a) + \frac{31}{18} \ln^2(a) \right. \\
\left. - \frac{4}{9} \ln^3(a) + 4 \sum_{n=1}^{\infty} \left[\frac{a^{-(n+1)}}{n(n+1)^3} - \frac{2a^{-(n+2)}}{n(n+2)^3} + \frac{a^{-(n+3)}}{n(n+3)^3} \right] \right\}. \tag{129}
\end{aligned}$$

Acknowledgments

This work was partially supported by NSF grant PHY-0244714 and by the Institute for Fundamental Theory.

-
- [1] R. P. Woodard, “Quantum effects during inflation,” *astro-ph/0310757*.
 - [2] V. K. Onemli and R. P. Woodard, *Class. Quant. Grav.* **19**, 4607 (2002), *gr-qc/0204065*.
 - [3] V. K. Onemli and R. P. Woodard, “Quantum effects can render $w < -1$ on cosmological scales,” *gr-qc/0406098*.
 - [4] A. Vilenkin and L. H. Ford, *Phys. Rev.* **D26**, 1231 (1982).

- [5] A. D. Linde, Phys. Lett. **116B**, 335 (1982).
- [6] A. A. Starobinskiĭ, Phys. Lett. **117B**, 175 (1982).
- [7] U. Alam, V. Sahni, T. D. Saini and A. A. Starobinskiĭ, “Is there evidence for dark energy metamorphosis?,” *astro-ph/0311364*.
- [8] T. R. Choudhury and T. Padmanabhan, “Cosmological parameters from supernova observations: A critical comparison of three data sets,” *astro-ph/0311622*.
- [9] Y. Wang and P. Mukherjee, Astrophys. J. **606**, 654 (2004), *astro-ph/0312192*.
- [10] Y. Gong, “Observational constraints on dark energy model,” *astro-ph/0401207*.
- [11] S. Nesseris and L. Perivolaropoulos, “A comparison of cosmological models using recent supernova data,” *astro-ph/0401556*.
- [12] Y. Wang and M. Tegmark, “New dark energy constraints from supernova, microwave background and galaxy clustering,” *astro-ph/0403292*.
- [13] U. Alam, V. Sahni and A. A. Starobinskiĭ, “The case for dynamical dark energy revisited,” *astro-ph/0403687*.
- [14] B. Feng, X. Wang and X. Zhang, “Dark energy constraints from the cosmic age and supernova,” *astro-ph/0404224*.
- [15] H. K. Jassal, J. S. Bagla and T. Padmanabhan, “WMAP constraints on low redshift evolution of dark energy,” *astro-ph/0404378*.
- [16] P. S. Corasaniti, M. Kunz, D. Parkinson, E. J. Copeland and B. A. Bassett, “The foundations of observing dark energy dynamics with the Wilkinson Microwave Anisotropy Probe,” *astro-ph/0406608*.
- [17] S. Hannestad and E. Mortsell, “Cosmological constraints on the dark energy equation of state and its evolution,” *astro-ph/0407259*.
- [18] J.-M. Virey, A. Ealet, C. Tao, A. Tilquin, A. Bonissent, D. Fouchez and P. Taxil, “Probing Dark Energy with Supernovae: a concordant or a convergent model?,” *astro-ph/0407452*.
- [19] S. M. Carroll, M. Hoffman and M. Trodden, Phys. Rev. **D68**, 023509 (2003), *astro-ph/0301273*.
- [20] J. M. Cline, S. Y. Jeon and G. D. Moore, “The phantom menaced: Constraints on low-energy effective ghosts,” *hep-ph/0311312*.
- [21] S. Nojiri, S. D. Odintsov and S. Ogushi, Int. J. Mod. Phys. **A17**, 4809 (2002), *hep-th/0205187*.
- [22] S. Nojiri and S. D. Odintsov, JCAP **0306**, 004 (2003), *hep-th/0303011*.

- [23] S. Nojiri and S. D. Odintsov, Phys. Lett. **562B**, 147 (2003), *hep-th/0303117*.
- [24] S. Nojiri and S. D. Odintsov, Phys. Lett. **565B**, 1 (2003), *hep-th/0304131*.
- [25] A. O. Barvinsky , Phys. Lett. **572B**, 109 (2003), *hep-th/0304229*.
- [26] S. Nojiri and S. D. Odintsov, Phys. Lett. **571B**, 1 (2003), *hep-th/0306212*.
- [27] S. Nojiri and S. D. Odintsov, “Gravity assisted dark energy dominance and cosmic acceleration,” *astro-ph/0403622*.
- [28] S. K. Srivastava, “Dark energy and dark matter of the universe from one-loop renormalization of riccion,” *hep-th/0404170*.
- [29] E. Elizalde, S. Nojiri and S. D. Odintsov, “Late-time cosmology in (phantom) scalar-tensor theory: Dark energy and the cosmic speed-up,” *hep-th/0405034*.
- [30] S. Nojiri and S. D. Odintsov, “The final state and thermodynamics of dark energy universe,” *hep-th/0408170*.
- [31] J. A. S. Lima and J. S. Alcaniz, “Thermodynamics and spectral distribution of dark energy,” *astro-ph/0402265*.
- [32] M. D. Maia, E. M. Monte, J. M. F. Maia and J. S. Alcaniz, “On the geometry of dark energy,” *astro-ph/0403072*.
- [33] J. S. Alcaniz and N. Pires, “Cosmic acceleration in brane cosmology,” *astro-ph/0404146*.
- [34] M. Bouhmadi-Lopez and J. Madrid, “Escaping the big rip?,” *astro-ph/0404540*.
- [35] P. H. Frampton and T. Takahasi, “Bigger rip with no dark energy,” *astro-ph/0405333*.
- [36] Y.-H. Wei, “The power-law expansion universe,” *astro-ph/0405368*.
- [37] S. D. H. Hsu, A. Jenkins and M. B. Wise, “Gradient instability for $w < -1$,” *astro-ph/0406043*.
- [38] S. K. Srivastava, “Future universe with $w < -1$ without big smash,” *astro-ph/0407048*.
- [39] A. Vikman, “Can dark energy evolve to the phantom?,” *astro-ph/0407107*.
- [40] E. Babichev, V. Dokuchaev and Y. Eroshenko, “Dark energy cosmology with generalized linear equation of state,” *astro-ph/0407190*.
- [41] S. Nojiri, “Dark energy and modified gravities,” *hep-th/0407099*.
- [42] B. Feng, M. Li, Y.-S. Piao and X. Zhang, “Oscillating quintom and the recurrent universe,” *astro-ph/0407432*.
- [43] L. Amendola, M. Gasperini and F. Piazza, “Fitting type Ia supernovae with coupled dark energy,” *astro-ph/0407573*.
- [44] F. C. Carvalho and A. Saa, “Non-minimal coupling, exponential potentials and the $w < -1$

- regime of dark energy,” *astro-ph/0408013*.
- [45] S. M. Carroll, A. De Felice and M. Trodden, “Can we be tricked into thinking that w is less than -1 ?” *astro-ph/0408081*.
 - [46] R. Holman and S. Naidu, “Dark energy from ‘water’,” *astro-ph/0408102*.
 - [47] S. Capozziello, A. Melchiorri and A. Schirone, “Scaling dark energy,” *astro-ph/0408112*.
 - [48] O. Bertolami and J. Paramos, “Astrophysical constraints on scalar field models,” *astro-ph/0408216*.
 - [49] Y.-H. Wei, “The power-law expansion universe and the late-time behavior,” *astro-ph/0408230*.
 - [50] Y.-H. Wei, “Late-time phantom universe in $SO(1,1)$ dark energy model with exponential potential,” *gr-qc/0410050*.
 - [51] A. Lue and G. D. Starkman, “How a brane cosmological constant can trick us into thinking that $w < -1$,” *astro-ph/0408246*.
 - [52] P. Wang and X. Meng, “Can vacuum decay in our universe,” *astro-ph/0408495*.
 - [53] A. O. Barvinsky, “Cosmological constant problem and long-distance modifications of Einstein theory,” *hep-th/0409144*.
 - [54] S. K. Srivastava, “Tachyon as a dark energy source,” *gr-qc/0409074*.
 - [55] T. Roman, “Some thoughts on energy conditions and wormholes,” *gr-qc/0409090*.
 - [56] W. Fang, H. Q. Lu, Z. G. Huang and K. F. Zhang, “Phantom cosmology with the Born-Infeld type scalar field,” *hep-th/0409080*.
 - [57] D. Boyanovski, H. J. de Vega and N. G. Sanchez, “Particle decay during inflation: self-decay of inflaton quantum fluctuations during slow roll,” *astro-ph/0409406*.
 - [58] D. Rapetti, S. W. Allen and J. Weller, “Constraining dark energy with X-ray galaxy clusters, supernovae and the cosmic microwave background,” *astro-ph/0409574*.
 - [59] C. Csaki, N. Kaloper and J. Terning, “Exorcising $w < -1$,” *astro-ph/0409596*.
 - [60] E. S. Santini and G. A. Lemarchand, “Accelerated expansion in a stochastic self-similar fractal universe,” *astro-ph/0410056*.
 - [61] L. P. Chimento and R. Lazkoz, “Large-scale inhomogeneities in modified Chaplygin gas cosmologies,” *astro-ph/0411068*.
 - [62] R. Lazkoz, “Rigidity of cosmic acceleration in a class of k-essence cosmologies,” *gr-qc/0410019*.
 - [63] L. P. Chimento and R. Lazkoz, “Unified Phantom Cosmologies,” *astro-ph/0405518*.
 - [64] J. M. Aguirregabiria, L. P. Chimento and R. Lazkoz, Phys. Rev. **D70**, 023509 (2004), *astro-*

ph/0403157.

- [65] L. P. Chimento and R. Lazkoz, Phys. Rev. Lett. **91**, 211301 (2003), *gr-qc/0307111*.
- [66] G. Calcagni, “Patch dualities and remarks on nonstandard cosmologies,” *gr-qc/0410027*.
- [67] J. Martin and M. A. Musso, “Stochastic Quintessence,” *astro-ph/0410190*.
- [68] S. Nesseris and L. Perivolaropoulos, “The fate of bound systems in phantom and quintessence cosmologies,” *astro-ph/0410309*.
- [69] R. J. Scherrer, “Phantom dark energy, cosmic doomsday, and the coincidence problem,” *astro-ph/0410508*.
- [70] E. Majerotto, D. Sapone and L. Amendola, “Supernovae type Ia data favour negatively coupled phantom energy,” *astro-ph/0410543*.
- [71] W. Hu, “Crossing the phantom divide: dark energy internal degrees of freedom,” *astro-ph/0410680*.
- [72] Z.-K. Guo, Y.-S. Piao, X.-M. Zhang and Y.-Z. Zhang, “Cosmological evolution of a quintom model of dark energy,” *astro-ph/0410654*.
- [73] P. P. Avelino, “The coincidence problem in linear dark energy models,” *astro-ph/0411033*.
- [74] T. Padmanabhan, “Dark energy: the cosmological challenge of the millennium,” *astro-ph/0411044*.
- [75] R.-G. Cai and A. Wang, “Cosmology with interaction between phantom dark energy and dark matter and the coincidence problem,” *hep-th/0411025*.
- [76] G. Huey, “A comprehensive approach to resolving the nature of the dark energy,” *astro-ph/0411102*.
- [77] D. A. Easson, “Cosmic acceleration and modified gravitational models,” *astro-ph/0411209*.
- [78] Y. Gong, “Observational constraints on generalized Chaplygin gas model,” *astro-ph/0411253*.
- [79] S. DeDeo, “Particle dark energy,” *astro-ph/0411283*.
- [80] J.-Q. Xia, B. Feng and X. Zhang, “Constraints on oscillating quintom from supernova, microwave background and galaxy clustering,” *astro-ph/0411501*.
- [81] Z.-K. Guo and Y.-Z. Zhang, “Interacting Phantom Energy,” *astro-ph/0411524*.
- [82] S. K. Srivastava, “Tachyon driven solution to cosmic coincidence problem,” *gr-qc/0411088*.
- [83] E. J. Copeland, M. R. Garousi, M. Sami and S. Tsujikawa, “What is needed of a tachyon if it is to be the dark energy?,” *hep-th/0411192*.
- [84] J. Schwinger, J. Math. Phys. **2**, 407 (1961).

- [85] R. D. Jordan, Phys. Rev. **D33**, 444 (1986).
- [86] B. Allen and A. Folacci, Phys. Rev. **D35**, 3771 (1987).
- [87] T. Prokopec and R. P. Woodard, Ann. Phys. **312**, 1 (2004), *gr-qc/0310056*.
- [88] T. Prokopec and R. P. Woodard, JHEP **0310**, 059 (2003), *astro-ph/0309593*.
- [89] A. A. Starobinskiĭ and J. Yokoyama, Phys. Rev. **D50**, 6357 (1994), *astro-ph/9407016*.
- [90] A. A. Starobinskiĭ, “Stochastic de Sitter (inflationary) stage in the early universe,” in *Field Theory, Quantum Gravity and Strings*, 107-126 (1986) ed. by H. J. de Vega and N. Sanchez.